arXiv:0907.5339v1 [math.CT] 30 Jul 2009

A Model Structure on the Category of Small Categories for Coverings

Kohei Tanaka

Department of Mathmatical Sciences Shinshu University, Matsumoto, Japan

July 30, 2009

Abstract

We define a new model structure on the category of small categories, which is intimately related to the notion of coverings and fundamental groups of small categories. Fibrant objects in the model structure coincide with groupoids, and the fibrant replacement is the groupoidification.

Contents

1 Introduction **1**

2 The fundamental groups of small categories **3**

3 The 1-type model structure on Cat **7**

4 The coverings of small categories **12**

5 Fibrations and coverings **17**

6 The classification theorem of coverings of small categories **23**

7 Relations between the 1-type model category and other model categories **27**

1 Introduction

The category **Cat** of small categories has a couple of interesting model structures. One of them is introduced by Joyal and Tierney [JT91], [Rez00], which is Quillen adjoint to the Joyal type model structure [Joy08] on the category **SSets** of simplicial sets and the Anderson type model structure [And78], [CGT04], [Str00] on the category **Grpd** of groupoids. On the other hand, Thomason found another model structure [Tho80] which is Quillen equivalent to the Kan type model structure [Qui67],[Hov99] on **SSets** and the Quillen type model structure [Qui67], [Hov99] on the category **Spaces** of topological spaces. These model categories are related to each other by the following functors

$$\begin{tikzcd}
\mathbf{Grpd} \arrow[r, shift right, "i"'] & \mathbf{Cat} \arrow[l, shift right, "\pi"'] \arrow[r, shift right, "N"'] & \mathbf{SSets} \arrow[l, shift right, "c"'] \arrow[r, shift right, "|-|"'] & \mathbf{Spaces} \arrow[l, shift right, "S"']
\end{tikzcd}$$

where i, π, N, c, $|-|$ and S are the inclusion, the groupoidification, the nerve, the categorization, the realization and the singular simplicial set functor, respectively. In [Qui68], Quillen shows that Serre fibrations in **Spaces** are related to Kan fibrations in **SSets** by $|-|$ and S. Similarly, Gabriel and Zisman define coverings of simplicial sets, and show that coverings in **Spaces** are related to coverings in **SSets** by $|-|$ and S in

[GZ67]. On the other hand, May defines coverings of groupoids, and show that coverings in **Spaces** are related to coverings in **Grpd** by the fundamental groupoid functor in [May99], [GZ67].

Now, we consider coverings in **Cat** related to coverings in **SSets** and **Grpd** by the above functors. The aim of this article is to show that we can treat coverings of small categories in terms of model categories. We prove **Cat** has a model structure, under which coverings are fibrations with discrete fibers.

Theorem 1.1. *There is a model structure on the category of small categories with the following weak equivalences, fibrations, and cofibrations.*

1. *A morphism $f : C \longrightarrow D$ is a weak equivalence iff $f_* : \pi_n(C, x) \longrightarrow \pi_n(D, f(x))$ is an isomorphism for $n = 0, 1$.*
2. *A morphism is a cofibration iff it is injective on the set of objects.*
3. *A morphism is a fibration iff it has the right lift property for all trivial cofibrations (weak equivalence and cofibration).*

We call the above model structure the "1-type model structure". The 1-type model structure on **SSets** and **Spaces** is already described in [DP95]. We prove Theorem 1.1 comparing **Cat** with **SSets** equipped with the 1-type model category. Furthermore, the model category in Theorem 1.1 has the following properties.

Theorem 1.2. *The category of small categories equipped with the* 1*-type model structure satisfies the following properties.*

1. *This is a cofibrantly generated model category.*
2. *This is Quillen equivalent to the category of groupoids with the Anderson type model structure in [And78], [Str00], [CGT04], and Quillen equivalent to the category of simplicial sets with the* 1*-type model structure [DP95].*
3. *Fibrant objects coincide with groupoids.*
4. *Coverings in the category of small categories coincide with fibrations with discrete fibers.*
5. *p is a fibration iff $N(p)$ is a Kan fibration.*
6. *The functorial factorization of the model category for the inclusion functor $* \longrightarrow C$ induces the universal cover over C.*

The paper is organized as follows. In section 2, we study the fundamental groups of small categories. The fundamental group is defined in terms of the "hammock localization" of a small category C introduced by Dwyer and Kan [DK80]. It coincides with the fundamental group defined by Minian in [Min02] and also isomorphic to $\pi_1(BC)$, where BC is the classifying space of C. In section 3, we prove theorem 1.1. In section 4, we define coverings in the category of small categories. The notion of coverings was already defined in the category of spaces, simplicial sets [GZ67], groupoids [May99], and k-linear categories [Gab81], [CRS07]. We consider the relations between them. And also we see the relations between coverings and fibrations in the 1-type model structure in section 5, between coverings and the fundamental groups of small categories in section 6. We prove the classification theorem of coverings in the category of small categories. This is a generalization of the classification theorem of coverings in the category of groupoids in [May99]. In section 7, we see the relations between the 1-type model category and other model categories.

Acknowledgement

I would like to thank Prof. Dai Tamaki and Katsuhiko Kuribayashi for helpful advices about this paper. Their several suggestions and remarks were essential in the development and revisions of this work. I am also grateful to Prof. Hideto Asashiba for conversations. His view points and suggestions really helped me.

2 The fundamental groups of small categories

Minian defined the fundamental group $\pi_1(C)$ of a small category C as the colimit of strong homotopy classes of functors from interval categories I_n to C for $n \geqq 0$ [Min02]. He also shows that $\pi_1(C) \cong \pi_1(BC)$.

Now, we define the fundamental group of a small category C as the endmorphism group of the groupoidification of C. We choose a model of groupoidification as $\pi_0(L^H C)$ where $L^H C$ is the hammock localization of C defined by Dwyer and Kan [DK80].

Definition 2.1. Given $n \in \mathbb{N}$, let I_n be the following category. The objects of I_n are the integers $0, 1, \cdots, n$ and the morphisms, other than the identities, are defined as follows. If r and s are two distinct objects in I_n there is exactly one morphism from r to s if r is even and $s = r - 1$ or $s = r + 1$ and no morphism otherwise. A sketch of I_n is as follow (case n even).

$$I_n : 0 \longrightarrow 1 \longleftarrow 2 \longrightarrow \cdots \longleftarrow n$$

Let C be a small category, denote the set of objects of C by C_0. Define a set $\pi_0(C) = C_0/\sim$ where $a \sim b$ iff there exists a functor $\alpha : I_n \longrightarrow C$ such that $\alpha(0) = a$ and $\alpha(n) = b$ for some n. $\pi_0(C)$ is called the set of path component of C.

Dwyer and Kan defined a simplicial category $L^H(C, W)$ with the same objects as C for a small category C and a subcategory W of C in [DK80], where the simplicial category means the category enriched by the category of simplicial sets. When $W = C$, we denote $L^H(C, W)$ as $L^H(C)$. For $a, b \in C_0$, k-simplices of the simplicial set $\mathrm{Hom}_{L^H(C)}(a, b)$ are called hammocks of width k. A hammock a diagram of the following form

$$\begin{tikzcd}
& C_{0,1} \arrow[r, dash] \arrow[d] & C_{0,2} \arrow[r, dash] \arrow[d] & \cdots \arrow[r, dash] & C_{0,n-1} \arrow[d] \arrow[ddr, dash] & \\
& C_{1,1} \arrow[r, dash] \arrow[d] & C_{1,2} \arrow[r, dash] \arrow[d] & \cdots \arrow[r, dash] & C_{1,n-1} \arrow[d] \arrow[dr, dash] & \\
X \arrow[uur, dash] \arrow[ur, dash] \arrow[r, dash] \arrow[dr, dash] \arrow[ddr, dash] & C_{2,1} \arrow[r, dash] \arrow[d, dotted, dash] & C_{2,2} \arrow[r, dash] \arrow[d, dotted, dash] & \cdots \arrow[r, dash] & C_{2,n-1} \arrow[r, dash] \arrow[d, dotted, dash] & Y \\
& C_{k-1,1} \arrow[r, dash] \arrow[d] & C_{k-1,2} \arrow[r, dash] \arrow[d] & \cdots \arrow[r, dash] & C_{k-1,n-1} \arrow[d] \arrow[ur, dash] & \\
& C_{k,1} \arrow[r, dash] & C_{k,2} \arrow[r, dash] & \cdots \arrow[r, dash] & C_{k,n-1} \arrow[uur, dash] &
\end{tikzcd}$$

for $n \geqq 0$, where the horizontal sequence means a functor $I_n \longrightarrow C$. The category $\pi_0(L^H(C))$ is a groupoid from Proposition 3.1 in [DK80]. We call $\pi_0(L^H(C))$ the groupoidification of C.

Definition 2.2. Define a set

$$(C^I; a, b) = \{\alpha : I_n \longrightarrow C \mid \alpha(0) = a, \alpha(n) = b, n \geqq 0\}$$

for $a, b \in C_0$, and we define a relation $\sim$ on $(C^I; a, b)$ as follow.

1. $(c \xrightarrow{f} d \xleftarrow{f} c) \sim c \sim (c \xleftarrow{f} d \xrightarrow{f} c)$
2. $(c \xrightarrow{f} d \xleftarrow{=} d \xrightarrow{g} e) \sim (c \xrightarrow{g \circ f} e)$, $(c \xleftarrow{f} d \xrightarrow{=} d \xleftarrow{g} e) \sim (c \xleftarrow{f \circ g} e)$
3. $(c \xrightarrow{f} b \xleftarrow{=} b) \sim (c \xrightarrow{f} b)$

$\pi(C; a, b)$ is defined by $(C^I; a, b)/\sim$.

The small category $\pi(C)$ is defined as $\pi(C)_0 = C_0$ and $\mathrm{Hom}_{\pi(C)}(a, b) = \pi(C; a, b)$. The composition is given by concatenation. Then every morphisms are invertible, so $\pi(C)$ is a groupoid and $\pi(C) = \pi_0(L^H(C))$. $\pi(C)$ gives a functor $\pi : \mathbf{Cat} \longrightarrow \mathbf{Grpd}$.

Define a functor $C \longrightarrow \pi(C)$ by $c \mapsto c$ on the set of objects and by $f \mapsto (f)$ on the set of morphisms, where (f) is the class of f.

For a pointed small category $(C,*)$, define a group $\pi_1(C,*)$ by $\mathrm{Hom}_{\pi(C)}(*,*)$. We call that C is path connected if $\pi_0(C) = *$ and simply connected if $\pi_0(C) = *$ and $\pi_1(C) = 1$.

Proposition 2.3. *π is a left adjoint functor to the inclusion functor $i : \mathbf{Grpd} \longrightarrow \mathbf{Cat}$.*

Proof. Let C be a small category and G be a groupoid. The functor $C \longrightarrow \pi(C)$ in Definition 2.2 induces a natural isomorphism

$$\mathrm{Hom}_{\mathbf{Grpd}}(\pi(C), G) \longrightarrow \mathrm{Hom}_{\mathbf{Cat}}(C, G)$$

then π is a left adjoint to i. □

π is called the groupoidification functor, and $\pi_1(C,*)$ is the endmorphism group of it on the object $*$. Next we study the construction of fundamental group given by Minian [Min02].

Definition 2.4. Let $(C,*)$ be a pointed category and let $f, g : I_n \longrightarrow C$ be functors such that $f(0) = f(n) = g(0) = g(n) = *$ for $n \geqq 0$. f and g are called strongly homotopic if there exists a functor $H : I_n \times I_m \longrightarrow C$ such that $H_{C\times\{0\}} = f$ and $H_{C\times\{m\}} = g$ and $H(0,k) = H(n,k) = *$, $(0 \leqq k \leqq m)$ for some $m \geqq 0$. Denote $f \simeq g$ when f and g are strongly homotopic.

$$[I_n, C]_* := \{f : I_n \longrightarrow C \mid f(0) = f(n) = *\}/\simeq$$

For $n > m$, define a functor $I_n \longrightarrow I_m$ by $i \mapsto i$ for $0 \leqq i \leqq m$ and $i \mapsto m$ for $i > m$.

$$\widetilde{\pi}_1(C,*) := \mathrm{colim}_n [I_n, C]_*.$$

The group structure on $\widetilde{\pi}_1(C,*)$ is defined by concatenation.

Proposition 2.5. *$\pi_1(C,*) = \widetilde{\pi}_1(C,*)$ for any pointed small category $(C,*)$.*

Proof. Both $\pi_1(C,*)$ and $\widetilde{\pi}_1(C,*)$ are consist of classes of zigzag sequences of C. See the each equivalence relations. The relation 3 in Definition 2.2 of $\pi_1(C)$ corresponds to the colimit relation of $\widetilde{\pi}_1(C,*)$. Let

$$* = c_0 \longrightarrow c_1 \longleftarrow c_2 \longrightarrow \cdots \longleftarrow c_n = *$$

and

$$* = d_0 \longrightarrow c_1 \longleftarrow c_2 \longrightarrow \cdots \longleftarrow d_n = *$$

be strongly homotopy equivalent. Then there exists a diagram

$$\begin{tikzcd}
\ast \arrow[r] \arrow[d, no head, "="'] & c_1 \arrow[d] & c_2 \arrow[l] \arrow[r] \arrow[d] & \cdots & c_{n-1} \arrow[l] \arrow[r] \arrow[d] & \ast \arrow[d, no head, "="] \\
\ast \arrow[r] \arrow[d, no head, "="'] & \cdot & \cdot \arrow[l] \arrow[r] & \cdots & \cdot \arrow[l] \arrow[r] & \ast \arrow[d, no head, "="] \\
\ast \arrow[r] \arrow[d, dotted, no head] & \cdot \arrow[u] \arrow[d, dotted, no head] & \cdot \arrow[l] \arrow[r] \arrow[u] \arrow[d, dotted, no head] & \cdots & \cdot \arrow[l] \arrow[r] \arrow[u] \arrow[d, dotted, no head] & \ast \arrow[d, dotted, no head] \\
\ast \arrow[r] \arrow[d, no head, "="'] & \cdot & \cdot \arrow[l] \arrow[r] & \cdots & \cdot \arrow[l] \arrow[r] & \ast \arrow[d, no head, "="] \\
\ast \arrow[r] & d_1 \arrow[u] & d_2 \arrow[l] \arrow[r] \arrow[u] & \cdots & d_{n-1} \arrow[l] \arrow[r] \arrow[u] & \ast
\end{tikzcd}$$

By chasing this diagram, we obtain

$$(* = c_0 \longrightarrow c_1 \longleftarrow c_2 \longrightarrow \cdots \longleftarrow c_n = *) = (* = d_0 \longrightarrow d_1 \longleftarrow d_2 \longrightarrow \cdots \longleftarrow d_n = *)$$

in $\pi_1(C,*)$. Conversely,

$$\begin{tikzcd}
c \arrow[r, "="] \arrow[d, "="'] & c \arrow[d, "="'] & c \arrow[l, "="'] \arrow[d, "="] \\
c \arrow[r, "f"'] & d & c \arrow[l, "f"]
\end{tikzcd}$$

is a commutative diagram, and the above two horizontal sequences are strongly homotopy equivalent. Similarly,

$$\begin{tikzcd}
c \arrow[r, "f"] \arrow[d, "="'] & d \arrow[d, "g"'] & d \arrow[l, "="'] \arrow[r, "g"] \arrow[d, "g"] & e \arrow[d, "="] \\
c \arrow[r, "g\circ f"'] & e & e \arrow[l, "="] \arrow[r, "="'] & e
\end{tikzcd}$$

is a commutative diagram, and the above two horizontal sequences are strongly homotopy equivalent. Then the relations on $\pi_1(C,*)$ and $\widetilde{\pi}_1(C,*)$ are equal and also group structures, thus $\pi_1(C,*) = \widetilde{\pi}_1(C,*)$. $\square$

Theorem 2.6. [Min02] *There exists a natural isomorphism $\widetilde{\pi}_1(C,*) \cong \pi_1(BC,*)$ for any pointed small category $(C,*)$, where BC is the classifying space of C.*

By the above theorem, $\pi_1(C,*)$ can be calculated in the category of spaces. However, we calculate it using the only method of small categories.

Proposition 2.7. *If a base point of C is an initial or terminal object, then $\pi_1(C,*) = 1$*

Proof. Let $*$ be an initial object and consider a sequence

$$(* = c_0 \xrightarrow{f_1} c_1 \xleftarrow{f_2} c_2 \xrightarrow{f_2} \cdots \longleftarrow c_n = *)$$

then there exists a unique morphism $\alpha_2 : * = c_0 \longrightarrow c_2$. On the other hand, since $\mathrm{Hom}(c_0, c_1)$ consists of a single point $f_2 \circ \alpha_2 = f_1$. Therefore,

$$\begin{aligned}
&(* = c_0 \xrightarrow{\alpha_2} c_2 \xleftarrow{=} c_2 \xrightarrow{f_2} c_1 \xleftarrow{f_2} c_2 \longrightarrow \cdots \longleftarrow c_n = *)\\
=&(* = c_0 \xrightarrow{f_3\circ\alpha_2} c_3 \longleftarrow \cdots \longleftarrow c_n = *).
\end{aligned}$$

By continuing this operation, the above sequence can be shown to be equivalent to $* \xrightarrow{=} *$, thus $\pi_1(C,*) = 1$. Similarly, we can prove that $\pi_1(C,*) = 1$ if $*$ is a terminal object. $\square$

Lemma 2.8. *Suppose C is a small category and $f : x \longrightarrow y$ is a morphism in C. Define a map*

$$f_* : \pi_1(C,x) \longrightarrow \pi_1(C,y)$$

by

$$(x \longrightarrow \cdot \longleftarrow \cdot \longrightarrow \cdots \longleftarrow x) \mapsto (y \xrightarrow{=} y \xleftarrow{f} x \longrightarrow \cdot \longleftarrow \cdot \longrightarrow \cdots \longleftarrow x \xrightarrow{f} y),$$

then f_* is an isomorphism.

Proof. An inverse map $\pi_1(C,y) \longrightarrow \pi_1(C,x)$ is defined by

$$(y \xrightarrow{g} \cdot \longleftarrow \cdot \longrightarrow \cdots \longleftarrow y) \mapsto (x \xrightarrow{g\circ f} \cdot \longleftarrow \cdot \longrightarrow \cdots \longleftarrow y \xrightarrow{=} y \xleftarrow{f} x).$$

$\square$

Lemma 2.9. *If C is path connected, then $\pi_1(C,c) \cong \pi_1(C,d)$ for all $c, d \in C_0$.*

Proof. By the assumption, there exists a zigzag sequence

$$c \longrightarrow \cdot \longleftarrow \cdot \longrightarrow \cdots \longleftarrow d.$$

From Lemma 2.8, we obtain an isomorphism $\pi_1(C, c) \longrightarrow \pi_1(C, d)$. ☐

We write $\pi_1(C, *)$ simply as $\pi_1(C)$, if C is path connected.

Example 2.10. Let S^1 be the category consisting of two objects $0, 1$ and two parallel morphisms f, g and identity morphisms,

$$\begin{tikzcd} 0 \arrow[r, shift left, "f"] \arrow[r, shift right, "g"'] & 1. \end{tikzcd}$$

$\pi_1(S^1)$ is generated by $(0 \xrightarrow{f} 1 \xleftarrow{g} 0)$, and $\pi_1(S^1) \cong \mathbb{Z}$.

Lemma 2.11. *Let C be a pointed category, and $(c \xrightarrow{f} d \xleftarrow{g} e) \in \mathrm{Hom}_{\pi(C)}(c, e)$. If g has an inverse, then*

$$(c \xrightarrow{f} d \xleftarrow{g} e) = (c \xrightarrow{g^{-1} \circ f} e)$$

in $\mathrm{Hom}_{\pi(C)}(c, e)$.

Proof. The diagram

$$\begin{tikzcd} c \arrow[r, "f"] \arrow[d, "="'] & d \arrow[d, "g^{-1}"'] & e \arrow[l, "g"'] \arrow[d, "="] \\ c \arrow[r, "g^{-1} \circ f"'] & e & e \arrow[l, "="] \end{tikzcd}$$

is commutative. ☐

Example 2.12. *Let G be a group regarded as a groupoid with one point set of objects. Then an element of $\pi_1(G)$ can be written as $(g_1, g_2, \cdots, g_n)$ where $g_i \in G$ for any $1 \leqq i \leqq n$. The lemma 2.11 implies that*

$$(g_1, g_2, \cdots, g_n) = \left(g_1 g_2^{-1} \cdots g_n^{(-1)^{n-1}}\right)$$

in $\pi_1(G)$. And we obtain an isomorphism $\pi_1(G) \longrightarrow G$ by

$$(g_1, g_2, \cdots, g_n) \mapsto g_1 g_2^{-1} \cdots g_n^{(-1)^{n-1}}.$$

Thus $\pi_1(G) \cong G$.

Corollary 2.13. *Let $(G, *)$ be a pointed groupoid. It is obvious that $\pi_1(G) \cong \pi_1(H)$ where H is a full subcategory of G with*

$$H_0 = \{g \in G_0 \mid [g] = [*] \in \pi_0(G)\}.$$

And H is equivalent to $\mathrm{Hom}_G(*, *)$ as categories. Example 2.12 implies that

$$\pi_1(G) \cong \mathrm{Hom}_G(*, *).$$

3 The 1-type model structure on Cat

Model categories, first introduced by Quillen in [Qui67], form the foundation of homotopy theory. This is a framework to do homotopy theory in general categories. In this section, we define a model structure on the category of small categories, which is closely related to coverings, fundamental groups and groupoidifications.

Definition 3.1. Suppose C is a category. A functorial factorization is an ordered pair (α, β) of functors $\mathrm{Mor}(C) \longrightarrow \mathrm{Mor}(C)$ such that $f = \beta(f) \circ \alpha(f)$ for all morphism f in C, where $\mathrm{Mor}(C)$ is the category of morphisms of M.

Definition 3.2. Let C be a category and $i : A \longrightarrow B$ and $p : X \longrightarrow Y$ be morphisms of C. Then we call that p has the right lifting property for i or i has the left lifting property for p if for every commutative diagram in C of the following form,

$$\begin{tikzcd}
A \arrow[r, "f"] \arrow[d, "i"'] & X \arrow[d, "p"] \\
B \arrow[r, "g"'] & Y
\end{tikzcd}$$

there is a morphism $h : B \longrightarrow X$ such that $h \circ i = f$ and $p \circ h = g$. If the above h exists uniquely, then we call that p has the unique right lifting property for i or i has the unique left lifting property for p.

Definition 3.3. Let C be a category closed under finite limits and colimits. If the following diagram in C

$$\begin{tikzcd}
A \arrow[r, "h"] \arrow[d, "f"'] & C \arrow[d, "g"] \\
B \arrow[r, "k"'] & D
\end{tikzcd}$$

is a pushout, then the morphism g is called the pushout of f along h. If the diagram is a pullback, then the morphism f is called the pullback of g along k.

Proposition 3.4. *Let C be a category closed under finite limits and colimits, and let p be a morphism in C.*

1. *The class of morphisms with the left lifting property for p is closed under pushout.*
2. *The class of morphisms with the right lifting property for p is closed under pullback.*

Proof. See in [Hir03]. □

Definition 3.5. [Hov99] A model structure on a category M consists of three subcategories of M called weak equivalences, cofibrations, and fibrations, and two functorial factorizations (α, β) and (γ, δ) satisfying the following properties.

1. If f and g are morphisms of M such that $g \circ f$ is defined and two of f, g and $g \circ f$ are weak equivalences, then so is third.
2. If f and g are morphisms of M such that f is a retract of g and g is weak equivalence, cofibration, or fibration then so is f.
3. Define a map to be a trivial cofibration if it is both a cofibration and a weak equivalence. Similarly, define a map to be a trivial fibration if it is both a fibration and a weak equivalence. Then trivial fibrations have the right lifting property for all cofibrations, and fibrations have the right lifting property for all trivial cofibrations.
4. For any morphism f in M, $\alpha(f)$ is a cofibration , $\beta(f)$ is a trivial fibration, $\gamma(f)$ is a trivial cofibration and $\delta(f)$ is a fibration.

A model category is a category M with all small limits and colimits together with a model structure on M.

It tends to be quite difficult to prove that a category admits a model structure. The above axioms are always hard to check. However, there exists a technique to prove that a category admits a good model structure called cofibrantly generated.

Definition 3.6. [Hov99], [Hir03] We say that a model category M is cofibrantly generated if there exists sets I and J of morphisms such that:

1. Both I and J permit the small objects argument [Hir03, 10.5.15].
2. Fibrations coincide with the morphisms which have the right lifting property for J.
3. Trivial fibrations coincide with the morphisms which have the right lifting property for I.

I is called a set of generating cofibrations, and J is called a set of generating trivial cofibrations.

Definition 3.7. We use the following notations.

1. ϕ is the empty category.
2. $*$ is the category with a unique object $*$ and the only identity morphism.
3. $[n]$ is the poset $0 < 1 < 2 < \cdots < n$ regarded as a category $0 \longrightarrow 1 \longrightarrow 2 \longrightarrow \cdots \longrightarrow n$.
4. S^0 is the coproduct of two $*$'s.
5. S^1 is the category with two point set of objects and has the parallel morphisms between them $0 \rightrightarrows 1$.
6. I_2 is the category $1 \longleftarrow 0 \longrightarrow 2$.
7. CS^1 is the category consisting of three point set of objects $0 \longrightarrow 1 \rightrightarrows 2$ where 0 is an initial object.
8. S^∞ is the simply connected groupoid with two point set of objects $0 \leftrightarrows 1$.

ϕ is a initial object and $*$ is a terminal object in **Cat**. We define a set of functors which are injective on set of objects

$$I = \{\phi \longrightarrow * \ , \ S^0 \longrightarrow [1] \ , \ S^1 \longrightarrow [1]\}.$$

Next, we define a functor $* \longrightarrow S^\infty$ by $* \mapsto 0$ on the set of objects, and $* \longrightarrow [1]$ by $* \mapsto 0$ on the set of objects, and let $I_2 \longrightarrow [2]$ and $CS^1 \longrightarrow [2]$ be functors with identity maps on the set of objects. Denote

$$K = \{* \longrightarrow S^\infty\} \ , \ J_1 = \{* \longrightarrow [1] \ , \ * \longrightarrow [1]^{op}\}$$

$$J_2 = \{I_2 \longrightarrow [2] \ , \ I_2^{op} \longrightarrow [2]^{op}\} \ , \ J_3 = \{CS^1 \longrightarrow [2] \ , \ (CS^1)^{op} \longrightarrow [2]^{op}\}$$

and $J = J_1 \cup J_2 \cup J_3$.

Theorem 3.8. [JT91],[Rez00] *There exists a model structure on* **Cat** *with the following weak equivalences, cofibrations, and fibrations.*

1. *A morphism is a weak equivalence iff it is an equivalence of categories.*
2. *A morphism is a cofibration iff it is injective on set of objects.*
3. *A morphism is a fibration iff it has the right lifting property for K.*

Furthermore, this is a cofibrantly generated model structure with generating cofibrations I and trivial cofibrations K.

This is called the Joyal-Tierney type model structure and denote the category of small categories equipped with this model structure by $\mathbf{Cat}_{JT}$. Also the category of groupoids **Grpd** has the restricted model structure of the above.

Theorem 3.9. [And78],[CGT04] *There exists a model structure on* **Grpd** *with the following weak equivalences, cofibrations, and fibrations.*

1. *A morphism is a weak equivalence iff it is an equivalence of categories.*
2. *A morphism is a cofibration iff it is injective on set of objects.*
3. *A morphism is a fibration iff it has the right lifting property for K.*

Thomason found another model structure on **Cat** in [Tho80] as follows .

Theorem 3.10. [Tho80] *There exists a cofibrantly generated model structure on* **Cat** *whose a weak equivalence f iff the classifying spaces Bf is a weak homotopy equivalence in* **Spaces***.*

In [Tho80], **Cat** with Thomason type model structure is closely related to **SSets** with Kan type model structure [Qui68], [Hov99].

Theorem 3.11. [Qui68], [Hov99] *There exists a model structure on* **SSets** *with the following weak equivalences, cofibrations, and fibrations.*

1. *A morphism f is a weak equivalence iff the geometric realization $|f|$ is a weak homotopy equivalence in* **Spaces***.*
2. *A morphism is a cofibration iff it is injective for all dimensions.*
3. *A morphism is a fibration iff it is a Kan fibration.*

Our aim in this section is to prove that the **Cat** has another model structure called the 1-type model structure.

Definition 3.12. A functor $f : C \longrightarrow D$ is called a weak 1-equivalence if the induced maps

$$f_* : \pi_0(C) \longrightarrow \pi_0(D)$$

and $f_* : \pi_1(C, x) \longrightarrow \pi_1(D, f(x))$ are both bijections for all $x \in C_0$.

Lemma 3.13. *Let G be a groupoid, then the canonical inclusion functor $G \longrightarrow \pi(G)$ is an isomorphism of categories.*

Proof. The inverse functor $\pi(G) \longrightarrow G$ is given by the identity map on the set of objects, and

$$(\cdot \xrightarrow{f_1} \cdot \xleftarrow{f_2} \cdot \xrightarrow{f_3} \cdots \xleftarrow{f_n} \cdot) \mapsto f_n^{-1} \circ \cdots \circ f_3 \circ f_2^{-1} \circ f_1$$

on the set of morphisms. □

Proposition 3.14. *The canonical inclusion functor $j : C \longrightarrow \pi(C)$ is a weak 1-equivalence for any small category C.*

Proof. Since $C_0 = \pi(C)_0$, the definition of $\pi(C)$ implies that the induced map $j_* : \pi_0(C) = \pi_0(\pi(C))$. By Lemma 3.13, j induces an isomorphism $\pi(C) \longrightarrow \pi(\pi(C))$ of categories, then $j_* : \pi_1(C, *) \longrightarrow \pi_1(\pi(C), *)$ is an isomorphism. □

Lemma 3.15. *A functor $f : C \longrightarrow D$ is a weak 1-equivalence iff $\pi(f) : \pi(C) \longrightarrow \pi(D)$ is an equivalence of categories.*

Proof. If $\pi(f)$ is an equivalence, then it is clear that f is a weak 1-equivalence by Proposition 3.14 and the following commutative diagram.

$$\begin{tikzcd}
C \arrow[r, "f"] \arrow[d, "j"'] & D \arrow[d, "j"] \\
\pi(C) \arrow[r, "\pi(f)"'] & \pi(D)
\end{tikzcd}$$

Let $f : C \longrightarrow D$ be a weak 1-equivalence. Since $\pi_n(BG) = 0$ for any groupoid G and $n \geqq 2$, then $\pi(f)_* : \pi_n(B\pi(C)) \longrightarrow \pi_n(B\pi(D))$ is an isomorphism for all $n \geqq 0$. This is a weak equivalence on **Cat** with Thomason type model structure in Theorem 3.10. However, if a functor is a weak equivalence in Thomason-type model structure between groupoids, then it is an equivalence of categories [CGT04]. Thus $\pi(f)$ is an equivalence of categories. □

Theorem 3.16. *There exists a model structure on* **Cat** *with the following weak equivalences, cofibrations, and fibrations.*

1. *A morphism is a weak equivalence iff it is a weak* 1*-equivalence.*
2. *A morphism is a cofibration iff it is an injection on set of objects.*
3. *A morphism is a fibration iff it has the right lifting property for J.*

Furthermore, this is a cofibrantly generated model structure with generating cofibrations I and trivial cofibrations J.

Denote the category of small categories equipped with the above model structure by $\mathbf{Cat}_1$. Cofibrations in $\mathbf{Cat}_1$ coincide with cofibrations in $\mathbf{Cat}_{JT}$.

It was already proved that **SSets** and **Spaces** admit model structures called the 1-type model structure in [DP95].

Definition 3.17. A morphism $f : X \longrightarrow Y$ in **Spaces** is called a weak 1-equivalence if the induced map $f_* : \pi_n(X, x) \longrightarrow \pi_n(Y, f(x))$ is an isomorphism for $n = 0, 1$ and $x \in X$. And a morphism f in **SSets** is called a weak 1-equivalence if $|f|$ is a weak 1-eqivalence in **Spaces**.

Remark 3.18. *By Theorem 2.6, f is a weak* 1*-equivalence in* **Cat** *iff Nf is a weak* 1*-equivalence in* **SSets**.

Theorem 3.19. [DP95] *There exists a model structure on* **SSets** *with the following weak equivalences, cofibrations, and fibrations.*

1. *A morphism is a weak equivalence iff it is a weak* 1*-equivalence.*
2. *A morphism is a fibration iff it has the right lifting property for J'.*
3. *A morphism is a cofibration iff it has the left lifting property for trivial fibrations.*

where

$$J' = \{\Lambda^n_j \longrightarrow \Delta[n], \Lambda^3_k \longrightarrow \partial\Delta[3] \mid 0 < n \leqq 2, 0 \leqq j \leqq n, 0 \leqq k \leqq 3\}$$

Furthermore, this is a cofibrantly generated model structure with generating cofibrations I' and trivial cofibrations J', where

$$I' = \{\partial\Delta[n] \longrightarrow \Delta[n] \mid 0 \leqq n \leqq 2\}.$$

We will prove Theorem 3.16 by using the following theorem, which is stated in more general form by Hirschhorn [Hir03] .

Theorem 3.20. *Let M be a cofibrantly generated model category with generating cofibrations A and generating trivial cofibrations B. Let C be a category that is closed under small limits and colimits, and let $F : M \Longleftrightarrow C : G$ be a pair of adjoint functors. If we let $FA = \{F(a) \mid a \in A\}$ and $FB = \{F(b) \mid b \in A\}$ and we define a weak equivalence in C as a morphism f which $G(f)$ is a weak equivalence in M, a fibration in C as a morphism has the right lifting property for FB, and a cofibration in C as a morphism has the left lifting property for trivial fibrations. If*

1. *both FA and FB permit the small objects argument [Hir03. 10.5.15] and*
2. *p is a weak equivalence in C if p has the left lifting property for fibrations in C.*

Then there exists a cofibrantly generated model structure on C, in which FA is a generating cofibrations and FB is a generating trivial cofibrations with the above weak equivalences, fibrations, and cofibrations.

The category of simplicial sets and the category of small categories are related by the nerve functor and categorization functor in [LTW79].

Definition 3.21. The nerve functor $N : \mathbf{Cat} \longrightarrow \mathbf{SSets}$ is defined by

$$N_n C = \mathrm{Hom}_{\mathbf{Cat}}([n], C)$$

and

$$d_i(f_1, \cdots, f_n) = (f_1, \cdots, f_{i-1}, f_{i+1} \circ f_i, f_{i+2}, \cdots, f_n)$$

and

$$s_j(f_1, \cdots, f_n) = (f_1, \cdots, f_j, 1, f_{j+1}, \cdots, f_n).$$

Next, the categorization functor $c : \mathbf{SSets} \longrightarrow \mathbf{Cat}$ is defined as follows. $cX_0 = X_0$ and the set of morphisms of cX consists of the classes of $x_1 x_2 \cdots x_n$, where $x_i \in X_1$ and $d_1 x_i = d_0 x_{i+1}$ in X_0 for $1 \leqq i \leqq n$ and $d_1(\sigma) \sim d_0(\sigma) d_2(\sigma)$ for $\sigma \in X_2$.

Proposition 3.22. [LTW79] $c : \mathbf{SSets} \Longleftrightarrow \mathbf{Cat} : N$ is an adjoint pair, and $cN \cong 1_{\mathbf{Cat}}$.

We will prove that Theorem 3.16 using Theorem 3.20.

Proof of Theorem 3.16. By Proposition 3.22, $c : \mathbf{SSets} \Longleftrightarrow \mathbf{Cat} : N$ is an adjoint pair, and Theorem 3.19, **SSets** admits a cofibrantly generated model structure with generating cofibrations I' and generating trivial cofibrations J'.

$$c(\partial\Delta[0] \longrightarrow \Delta[0]) = \phi \longrightarrow *$$

and

$$c(\partial\Delta[1] \longrightarrow \Delta[1]) = S^0 \longrightarrow [1].$$

Now $c\partial\Delta[2]$ is of the following form.

$$\begin{tikzcd}
0 \arrow[rr, bend left] \arrow[dr] & & 2 \\
& 1 \arrow[ur] &
\end{tikzcd}$$

then $S^1 \longrightarrow [1]$ is a retract of $c(\partial\Delta[2] \longrightarrow \Delta[2])$. Conversely, the pushout of the inclusion functor

$$S^1 \longrightarrow c(\partial\Delta[2])$$

along $S^1 \longrightarrow [1]$ in Definition 3.3 is $c(\partial\Delta[2] \longrightarrow \Delta[2])$. Thus, the morphisms in **Cat** which have the right lifting property for $c(I')$ coincide with the morphisms which have the right lifting property for I. Similarly, $c(J')$ consists of identity maps and the pushout of morphisms of J, and J consists of the morphisms which are retracts of $c(J')$. Thus, the morphisms in **Cat** which have the right lifting property for $c(J')$ coincide with the morphisms which have the right lifting property for J. Then, the fibrations in Theorem 3.20 coincides with fibrations in Theorem 3.16. Since any category of domain of FI' and FJ' consists of finite objects and finite morphisms, then condition 1 of Theorem 3.20 is satisfied.

We will show that condition 2 of Theorem 3.20 holds. Let $j : A \longrightarrow X$ be a morphism in **Cat** which has the left lifting property for fibrations. Consider the following diagram in **Grpd**,

$$\begin{tikzcd}
\pi(A) \arrow[r] \arrow[d, "\pi(j)"'] & E \arrow[d, "p"] \\
\pi(X) \arrow[r] & B
\end{tikzcd}$$

where p is a fibration in **Grpd**. By Proposition 2.3, $\pi : \mathbf{Cat} \Longleftrightarrow \mathbf{Grpd} : i$ is an adjoint pair, then the above diagram induces a diagram in **Cat** as follows.

$$\begin{tikzcd}
A \arrow[r] \arrow[d, "j"'] & E \arrow[d, "p"] \\
X \arrow[r] & B
\end{tikzcd}$$

Now, $p : E \longrightarrow B$ as a morphism of **Cat** is a fibration in Definition 3.16 since $\pi(J)$ consists of identity maps and $* \longrightarrow S^{\infty}$. Then there exists a morphism $k : X \longrightarrow E$ making the following diagram commutative.

$$\begin{tikzcd}
A \arrow[r] \arrow[d, "j"'] & E \arrow[d, "p"] \\
X \arrow[r] \arrow[ur, "k"] & B
\end{tikzcd}$$

Then

$$\begin{tikzcd}
\pi(A) \arrow[r] \arrow[d, "\pi(j)"'] & E \arrow[d, "p"] \\
\pi(X) \arrow[r] \arrow[ur, "k^{\sharp}"] & B
\end{tikzcd}$$

is commutative, thus $\pi(j) : \pi(A) \longrightarrow \pi(X)$ is a trivial cofibration in **Grpd**. In particular, $\pi(j)$ is an equivalence of categories, then Proposition 3.15 implies that j is a weak 1-equivalence in **Cat**, and $N(f)$ is a weak 1-equivalence in **SSets** from Remark 3.18. The result follows from Theorem 3.20. $\square$

Remark 3.23. *The model structure on* **Cat** *in Theorem 3.16 coincide with the model structure in Theorem 1.1, since fibrations in a model category coincide with the morphisms which has right lifting property for trivial cofibrations [Hov99].*

4 The coverings of small categories

The notion of coverings is already defined in the category of spaces, simplicial sets [GZ67], groupoids [May99] and k-linear categories [Gab81], [CRS07].

Now we define coverings in the category of small categories, and consider the relations between our coverings and coverings defined in other categories.

Definition 4.1. A functor $p : E \longrightarrow B$ is called a covering if it has the unique right lifting property for the inclusion functors $0 : * \longrightarrow [1]$ and $1 : * \longrightarrow [1]$. A covering $p : E \longrightarrow B$ is called universal if E is simply connected.

Lemma 4.2. *$p : E \longrightarrow B$ is a covering iff p has the unique right lifting property for the inclusion functors $0 : * \longrightarrow I_n$ for $n > 0$.*

Proof. Since $I_n = 0 \longrightarrow 1 \longleftarrow 2 \longrightarrow \cdots \longleftarrow n$, then we repeat taking lifts of $i \longrightarrow i+1$ or $i \longleftarrow i+1$. $\square$

Proposition 4.3. *Let $p : E \longrightarrow B$ be a covering, then it has the unique right lifting property for an inclusion functor $(0,0) : * \longrightarrow I_n \times I_m$ for all $n, m > 0$.*

Proof. When $n = 1, m = 1$, given a diagram

$$\begin{tikzcd}
* \arrow[r] \arrow[d, "{(0,0)}"'] & E \arrow[d, "p"] \\
{[1] \times [1]} \arrow[r] & B
\end{tikzcd}$$

The bottom morphism $[1] \times [1] \longrightarrow B$ gives a commutative diagram in B as follows.

$$\begin{tikzcd}
b_0 = p(e) \arrow[r, "f"] \arrow[d, "j"'] & b_1 \arrow[d, "k"] \\
b_2 \arrow[r, "g"'] & b_3
\end{tikzcd}$$

Since p is a covering, we obtain morphisms $\tilde{f} : e \longrightarrow e_1$ and $\tilde{j} : e \longrightarrow e_2$ in E such that $p(\tilde{f}) = f$ and $p(\tilde{j}) = j$. Furthermore, we obtain morphisms $\tilde{g} : e_1 \longrightarrow e_3$ and $\tilde{k} : e_2 \longrightarrow e_3'$ in E such that $p(\tilde{g}) = g$ and $p(\tilde{k}) = k$. Now $p(\tilde{k} \circ \tilde{f}) = p(\tilde{g} \circ \tilde{j})$ in B, then $\tilde{k} \circ \tilde{f} = \tilde{g} \circ \tilde{j}$ in E since the unique lifting property of p. The commutative diagram in E

$$\begin{tikzcd}
e \arrow[r, "\tilde{f}"] \arrow[d, "\tilde{j}"'] & e_1 \arrow[d, "\tilde{k}"] \\
e_2 \arrow[r, "\tilde{g}"'] & e_3
\end{tikzcd}$$

gives a functor $[1] \times [1] \longrightarrow E$ making the diagram

$$\begin{tikzcd}
* \arrow[r] \arrow[d, "{(0,0)}"'] & E \arrow[d, "p"] \\
{[1] \times [1]} \arrow[r] & B
\end{tikzcd}$$

commutative. In general $n, m > 0$, we repeat taking lifts of diagrams. ☐

Corollary 4.4. *Let $p : E \longrightarrow B$ be a covering and $e, e' \in E_0$, $f, g : I_n \longrightarrow B$,*

$$f(0) = g(0) = p(e), \; f(n) = g(n) = p(e')$$

and $H : I_n \times I_m \longrightarrow B$ be a strong homotopy from f to g such that $H(0, a) = p(e), H(n, a) = p(e')$ for all $0 \leqq a \leqq m$. Denote the lifted morphisms of f, g by $\tilde{f}, \tilde{g} : I_n \longrightarrow E$ satisfying $\tilde{f}(0) = \tilde{g}(0) = e$, then $\tilde{f}(n) = \tilde{g}(n) = e'$ and there exists a strong homotopy $\widetilde{H}$ from $\tilde{f}$ to $\tilde{g}$ such that $\widetilde{H}(0, a) = e, \widetilde{H}(n, a) = e'$ for all $0 \leqq a \leqq m$.

Proof. By Proposition 4.3, H can be lifted to E. And we obtain a strong homotopy we want because of the unique lifting property of p. ☐

Proposition 4.5. *Let $p : E \longrightarrow B$ be a covering and $b \in B_0$, $e \in p^{-1}(b)$. Then the induced map $p_* : \pi_1(E, e) \longrightarrow \pi_1(B, b)$ is injective.*

Proof. Assume that $p(e_*) = 1$ in $\pi_1(B, b)$ for $e_* \in \pi_1(E, e)$. Thus $p(e_*) \simeq b$. The Corollary 4.4 implies that $e_* \simeq e$, therefore $e_* = 1 \in \pi_1(E, e)$. ☐

Next we construct the universal cover over a path connected category.

Definition 4.6. The Grothendieck construction or the transportation category [Tho79] of a functor $F : I \longrightarrow$ **Sets**, where I is a small category and **Sets** is the category of sets, is a small category $Gr(F)$ defined as follows. The set of objects of $Gr(F)$ consists of (i, x) where i is an object of I and x is an element of $F(i)$. And a morphism $(i, x) \longrightarrow (j, y)$ in $Gr(F)$ is a morphism $f : i \longrightarrow j$ in I such that $F(f)(x) = y$. And the canonical projection functor $Gr(F) \longrightarrow I$ is given by $(i, x) \longrightarrow i$.

Let $(C, *)$ be a pointed category, then the set of morphisms of $\pi(C)$ gives a functor

$$\mathrm{Hom}_{\pi(C)}(*, -) : C \longrightarrow \mathbf{Set}.$$

Denote $Gr\left(\mathrm{Hom}_{\pi(C)}(*,-)\right)$ by $\widehat{C}$ and the canonical projection by $T : \widehat{C} \longrightarrow C$. That is, an object of $\widehat{C}$ can be appeared as a equivalent class of a sequence in C

$$(* \longrightarrow c_1 \longleftarrow c_2 \longrightarrow \cdots \longleftarrow c_n)$$

and T carries it to the last object c_n.

Lemma 4.7. *Let C be a path connected category, then $T : \widehat{C} \longrightarrow C$ is a covering.*

Proof. Given a commutative diagram

$$\begin{tikzcd}
* \arrow[r, "f"] \arrow[d, "0"'] & \widehat{C} \arrow[d, "T"] \\
{[1]} \arrow[r, "g"'] & C
\end{tikzcd}$$

then, $f(0)$ is a class of zigzag sequence of C and $g(0 \longrightarrow 1) : g(0) = T(f(0)) \longrightarrow g(1)$. Define $h : [1] \longrightarrow \widehat{C}$ by $h(0) = f(0)$, $h(1) = (g(0 \longrightarrow 1)) \circ f(0)$ and $h(0 \longrightarrow 1) = g(0 \longrightarrow 1)$. It is obvious that

$$\begin{tikzcd}
* \arrow[r, "f"] \arrow[d, "0"'] & \widehat{C} \arrow[d, "T"] \\
{[1]} \arrow[r, "g"'] \arrow[ur, "h"] & C
\end{tikzcd}$$

is commutative and h exists uniquely. Similarly, T has the unique lifting property for $1 : * \longrightarrow [1]$. ☐

Proposition 4.8. *$\pi_1(\widehat{C}) = 1$ if C is path connected.*

Proof. For $(*) = (c_*^0) \xrightarrow{f_1} (c_*^1) \xleftarrow{f_2} (c_*^2) \longrightarrow \cdots \xleftarrow{f_n} (c_*^n) = (*) \in \pi_1(\widehat{C})$, it suffices to show that

$$(* \xrightarrow{f_1} T(c_*^1) \xleftarrow{f_2} T(c_*^2) \longrightarrow \cdots \xleftarrow{f_n} *) = 0 \in \pi_1(C).$$

We continue to deform the above sequence,

$$\begin{aligned}
&(* \xrightarrow{f_1} T(c_*^1) \xleftarrow{f_2} T(c_*^2) \longrightarrow \cdots \xleftarrow{f_n} *) \\
=&(* \xrightarrow{c_*^1} T(c_*^1) \xleftarrow{f_2} T(c_*^2) \longrightarrow \cdots \xleftarrow{f_n} *) \\
=&(* \xrightarrow{c_*^2} T(c_*^2) \longrightarrow \cdots \xleftarrow{f_n} *) \\
=&\cdots = *
\end{aligned}$$

☐

Corollary 4.9. *$T : \widehat{C} \longrightarrow C$ in definition 4.6 is a universal cover if C is path connected.*

Proof. By Proposition 4.8 and Lemma 4.7. ☐

Proposition 4.10. *Let C and E be path connected and pointed small categories and $q : E \longrightarrow C$ be a covering preserving the base point, then there exists a unique functor $\alpha : \widehat{C} \longrightarrow E$ such that preserving the base point and $q \circ \alpha = T$ where $T : \widehat{C} \longrightarrow C$ is the universal cover over C in Definition 4.6.*

Proof. Let $c_* \in \widehat{C}_0$, then the following commutative diagram

$$\begin{tikzcd}
* \arrow[r, "*"] \arrow[d] & E \arrow[d, "q"] \\
I_n \arrow[r, "c_*"'] & C
\end{tikzcd}$$

gives a functor $\tilde{c}_* : I_n \longrightarrow E$ such that $\tilde{c}(0) = *$ and $q \circ \tilde{c}_* = c_*$ since q is a covering. Thus, we define a correspondence $\alpha : \widehat{C} \longrightarrow E$ on the set of objects as $\alpha(c_*) = \tilde{c}_n$. A morphism $f : c_* \longrightarrow d_*$ in $\widehat{C}$ is a morphism $f : c_n \longrightarrow d_m$ in C between last objects of c_* and d_*, then we obtain a morphism in E, $\tilde{f} : \tilde{c}_n \longrightarrow \tilde{d}_m$ such that $p(\tilde{f}) = f$ and $\tilde{f} \circ \tilde{c}_* = \tilde{d}_*$ in $\pi(E)$. Then define α on the set of morphism as $\alpha(f) = \tilde{f}$. α is a functor $\widehat{C} \longrightarrow E$ from the unique lifting property of q and $q \circ \alpha = T$, $\alpha(*) = *$.

Now, let $\beta : \widehat{C} \longrightarrow E$ be a functor preserving the base point and $q \circ \beta = p$. For an object of $\widehat{C}$

$$c_* = (* \xrightarrow{f_1} c_1 \xleftarrow{f_2} \cdots \xleftarrow{f_n} c_n)$$

induces a zigzag sequence of $\widehat{C}$ as follows.

$$f_* : (*) \xrightarrow{f_1} (* \xrightarrow{f_1} c_1) \xleftarrow{f_2} \cdots \xleftarrow{f_n} (c_*)$$

Then $\beta(f_*)$ is a zigzag sequence of E starting the base point of E. For the following commutative diagram,

$$\begin{tikzcd}
* \arrow[r, "*"] \arrow[d] & E \arrow[d, "q"] \\
I_n \arrow[r, "c_*"'] & C
\end{tikzcd}$$

$\beta(f_*) : I_n \longrightarrow E$ is a functor that $p \circ \beta(f_*) = c_*$ and $\beta(f_*)(0) = *$. Since the unique lifting property, $\beta(f_*) = \tilde{c_*}$. Then $\beta(c_*) = \beta(f_*)(n) = \tilde{c}_n = \alpha(c_*)$, thus $\alpha = \beta$. □

We recall the definition of coverings in the category of simplicial sets, groupoids, and k-linear categories.

Definition 4.11. [GZ67] A morphism in the category of simplicial sets $p : E \longrightarrow B$ is called a covering if it has the unique right lifting property for the inclusions $\Delta[0] \longrightarrow \Delta[n]$, $n \geqq 0$.

Definition 4.12. [May99] A morphism in the category of groupoids $p : E \longrightarrow B$ is called a covering if it has the unique right lifting property for the inclusion $* \longrightarrow S^\infty$ in Definition 3.7.

Definition 4.13. [Gab81], [CRS07] A morphism in the category of k-linear categories $p : E \longrightarrow B$ is called a covering if

$$p : \bigoplus_{x \in E_0} \mathrm{Hom}_E(x, e) \longrightarrow \bigoplus_{y \in B_0} \mathrm{Hom}_B(y, b)$$

and

$$p : \bigoplus_{x \in E_0} \mathrm{Hom}_E(e, x) \longrightarrow \bigoplus_{y \in B_0} \mathrm{Hom}_B(b, y)$$

are isomorphisms for any $b \in B_0$ and $e \in p^{-1}(b)$.

Next we recall the relations between coverings in those categories.

Proposition 4.14. [GZ67] *S : **Spaces** $\longrightarrow$ **SSets** and $|-|$: **SSets** $\longrightarrow$ **Spaces** preserve coverings.*

Proposition 4.15. *i : **Grpd** $\longrightarrow$ **Cat** and π : **Cat** $\longrightarrow$ **Grpd** preserve coverings.*

Proof. i is a right adjoint to π, then i preserves coverings since $\pi([1]) = S^{\infty}$. Conversely, let $p : E \longrightarrow B$ be a covering in **Cat**. Consider the following diagram,

$$\begin{tikzcd}
* \arrow[r] \arrow[d, "e"'] & \pi(E) \arrow[d, "\pi(p)"] \\
S^{\infty} \arrow[r, "f"'] & \pi(B)
\end{tikzcd}$$

then $f(0 \longrightarrow 1) : I_n \longrightarrow B$ and $f(0 \longrightarrow 1)(0) = p(e)$.

$$\begin{tikzcd}
* \arrow[r, "e"] \arrow[d, "0"'] & E \arrow[d, "p"] \\
I_n \arrow[r, "f(0\longrightarrow 1)"'] & B
\end{tikzcd}$$

is a commutative diagram, then there exists a morphism $g : I_n \longrightarrow E$ such that $p \circ g = f(0 \longrightarrow 1)$ and $g(0) = e$. Thus

$$\begin{tikzcd}
* \arrow[r] \arrow[d, "e"'] & \pi(E) \arrow[d, "\pi(p)"] \\
S^{\infty} \arrow[r, "f"'] \arrow[ur, "\tilde{g}"] & \pi(B)
\end{tikzcd}$$

is commutative, where $\tilde{g}(0) = e$ and $\tilde{g}(0 \longrightarrow 1) = g$. Therefore $\pi(p)$ is a covering in the category of groupoids. □

Proposition 4.16. *For $N : \mathbf{Cat} \longrightarrow \mathbf{SSets}$, p is a covering in* **Cat** *iff $N(p)$ is a covering in* **SSets**.

Proof. N is a right adjoint functor of c, then N preserves coverings. Conversely, let $N(p)$ be a covering for p, then $N(p)$ has the unique right lifting property for

$$d_0^*, d_1^* : \Delta[0] \longrightarrow \Delta[1].$$

Since $cN \cong 1_{\mathbf{Cat}}$, then p has the unique right lifting property for $0 : * \longrightarrow [1]$ and $1 : * \longrightarrow [1]$. □

Lemma 4.17. *$p : E \longrightarrow B$ is a covering in* **Cat** *iff*

$$p : \coprod_{x \in E_0} \mathrm{Hom}_E(e, x) \longrightarrow \coprod_{y \in B_0} \mathrm{Hom}_B(b, y)$$

and

$$p : \coprod_{x \in E_0} \mathrm{Hom}_E(x, e) \longrightarrow \coprod_{y \in B_0} \mathrm{Hom}_B(y, b)$$

are bijective.

Proof. It is obvious that p has the unique lifting property for $0 : * \longrightarrow [1]$ iff

$$p : \coprod_{x \in E_0} \mathrm{Hom}_E(e, x) \longrightarrow \coprod_{y \in B_0} \mathrm{Hom}_B(b, y)$$

is bijective for $b \in B_0$ and $e \in p^{-1}(b)$. And p has the unique lifting property for $1 : * \longrightarrow [1]$ iff

$$p : \coprod_{x \in E_0} \mathrm{Hom}_E(x, e) \longrightarrow \coprod_{y \in B_0} \mathrm{Hom}_B(y, b)$$

is bijective.. □

Definition 4.18. Let k be a ring. Define a functor $k : \mathbf{Cat} \longrightarrow k\mathbf{Cat}$ as follows, where $k\mathbf{Cat}$ is the category of k-linear categories. $k(C)_0 = C_0$ and $\mathrm{Hom}_{kC}(a,b) = k(\mathrm{Hom}_C(a,b))$, that is the free k-module generated by $\mathrm{Hom}_C(a,b)$.

Proposition 4.19. *$k : \mathbf{Cat} \longrightarrow k\mathbf{Cat}$ in Definition 4.18 preserves coverings.*

Proof. Let p be a covering in $\mathbf{Cat}$, since Lemma 4.17,

$$p : \coprod_{x \in E_0} \mathrm{Hom}_E(e,x) \longrightarrow \coprod_{y \in B_0} \mathrm{Hom}_B(b,y)$$

and

$$p : \coprod_{x \in E_0} \mathrm{Hom}_E(x,e) \longrightarrow \coprod_{y \in B_0} \mathrm{Hom}_B(y,b)$$

are isomorphisms. Thus

$$kp : \bigoplus_{x \in E_0} k(\mathrm{Hom}_E(e,x)) \longrightarrow \bigoplus_{y \in B_0} k(\mathrm{Hom}_B(b,y))$$

and

$$kp : \bigoplus_{x \in E_0} k(\mathrm{Hom}_E(x,e)) \longrightarrow \bigoplus_{y \in B_0} k(\mathrm{Hom}_B(y,b))$$

are isomorphisms as morphisms in the category of k-modules. Then kp is a covering in $k\mathbf{Cat}$. ☐

5 Fibrations and coverings

Next we investigate properties of the 1-type model structure. In particular, we show that fibrations in the 1-type model category are closely related to coverings, prefibered functors, and Kan fibrations.

Definition 5.1. Suppose M be a model category, then M has an initial object ϕ and a terminal object $*$. We call an object X of M cofibrant if $\phi \longrightarrow X$ is a cofibration, and we call an object Y of M fibrant if $Y \longrightarrow *$ is a fibration.

Proposition 5.2. *In the 1-type model structure on* $\mathbf{Cat}$*, all objects are cofibrant, and fibrant objects coincide with groupoids.*

Proof. By the definition of cofibration in theorem 3.16, it is obvious that all objects are cofibrant. Next, let G be a groupoid, then $G \longrightarrow *$ has the lift property for J since all morphisms of G are invertible. Conversely, Let C be a fibrant object. For any morphism $f \in \mathrm{Hom}(x,y)$, we can consider a diagram in C that $y \xleftarrow{f} x \xrightarrow{=} x$. Use the lifting property for J_2, we obtain a morphism $g : y \longrightarrow x$ satisfying $g \circ f = 1$. Now $g \circ (f \circ g) = g = g \circ 1$, the lift property for J_3 implies that $f \circ g = 1$. Thus C is a groupoid. ☐

Lemma 5.3. *If $p : E \longrightarrow B$ is a covering, then p is a fibration in $\mathbf{Cat}_1$.*

Proof. Since the definition of coverings in Definition 4.1, p has the lift property for J_1. Next, given the following commutative diagram.

$$\begin{tikzcd}
I_2 \arrow[r, "f"] \arrow[d] & E \arrow[d, "p"] \\
{[2]} \arrow[r, "g"'] & B
\end{tikzcd}$$

Then, we obtain a morphism $\alpha : f(1) \longrightarrow e$ over $g(1 \longrightarrow 2) : g(1) \longrightarrow g(2)$ by the lifting property of p. And $p(\alpha \circ f(0 \longrightarrow 1)) = g(0 \longrightarrow 2)$ implies that $\alpha \circ f(0 \longrightarrow 1) = f(0 \longrightarrow 2)$ by the universality of the lifting property, then p has the lifting property for J_2. At last, universality of the lifting property implies that p has the lifting property for J_3. ☐

Definition 5.4. [Qui68] Let $f : X \longrightarrow Y$ be a functor, then the category $f^{-1}(y)$ is defined as a subcategory of X for $y \in Y_0$, $f^{-1}(y)_0 = f^{-1}(y)$ and $\mathrm{Hom}_{f^{-1}(y)}(a, b) = p^{-1}(1_y)$.

Furthermore, let y/f denote the category as $(y/f)_0 = \{(x, w) \in X_0 \times \mathrm{Hom}_Y(y, f(x))\}$ and

$$\mathrm{Hom}_{y/f}\left((x, w), (x', w')\right) = \{g \in \mathrm{Hom}_X(x, x') \mid f(g) \circ w = w'\},$$

similarly f/y denote the category as $(f/y)_0 = \{(x, w) \in X_0 \times \mathrm{Hom}_Y(f(x), y)\}$ and

$$\mathrm{Hom}_{f/y}\left((x, w), (x', w')\right) = \{g \in \mathrm{Hom}_X(x, x') \mid w' \circ f(g) = w\}.$$

Define functors $i_y : f^{-1}(y) \longrightarrow y/f$ and $j_y : f^{-1}(y) \longrightarrow f/y$ as $x \mapsto (x, 1_y)$ on the set of objects. We call f is a prefibered category if i_y has a right adjoint functor for all y, and we call f is a precofibered category if j_y has a left adjoint functor for all y.

Definition 5.5. A category is called discrete if the set of morphisms consists of only identity maps.

Proposition 5.6. *$p : E \longrightarrow B$ is a covering iff p is a fibration in $\mathbf{Cat}_1$ and the category of fiber $p^{-1}(b)$ is discrete for any $b \in B_0$.*

Proof. Let $p : E \longrightarrow B$ be a covering, then p is a fibration by lemma 5.3, and fibers have only identity maps since the universality of lifting property. Conversely, let p be a fibration with discrete fibers. Since p is a fibration, p has the lifting property for J_1. We will show that the lifting property has the universality. For a following commutative diagram,

$$\begin{tikzcd}
* \arrow[r] \arrow[d] & E \arrow[d, "p"] \\
{[1]} \arrow[r, "f"'] & B
\end{tikzcd}$$

assume $g, h : [1] \longrightarrow E$ are both satisfy $p \circ g = p \circ h = f$ and $g(0) = h(0) = e$. The lifting property of p for J_2 implies that there exists $w : g(1) \longrightarrow h(1)$ such that $w \circ g = h$ and $p \circ w = 1$. Then w is a morphism of $p^{-1}(f(1))$. However, $p^{-1}(f(1))$ has only identity maps, thus $w = 1$. Therefore, $g = h$. □

Proposition 5.7. *Let $p : E \longrightarrow B$ be a fibration, then p is both a prefibered category and a precofibered category.*

Proof. For $b \in B_0$, we define a left adjoint functor $r : p/b \longrightarrow p^{-1}(b)$ of $i : p^{-1}(b) \longrightarrow p/b$ as follows. Take an object $(e, w) \in (b/p)_0$, then $w : p(e) \longrightarrow b$ in E. Consider the following commutative diagram.

$$\begin{tikzcd}
* \arrow[r, "e"] \arrow[d] & E \arrow[d, "p"] \\
{[1]} \arrow[r, "w"'] & B
\end{tikzcd}$$

There exists a functor $\tilde{w} : [1] \longrightarrow E$ such that $p \circ \tilde{w} = w$ and $\tilde{w}(0) = e$ because of the lifting property of q for J_1. Therefore, define on set of objects as $r(e, w) = \tilde{w}(1)$. Next, for a morphism $\alpha : (e, w) \longrightarrow (e', w')$ in p/b, then $\alpha : e \longrightarrow e'$ in E such that $w' \circ p(\alpha) = w$. Since p has the lifting property for $I_2 \longrightarrow [2]$, then there exists a morphism $\beta : \tilde{w}(1) \longrightarrow \tilde{w'}(1)$ such that $\tilde{w'} \circ \alpha = \beta \circ w$ and $p(\beta) = 1_b$. Thus, define $r(\alpha) = \beta$. Because of the lifting property of p for $CS^1 \longrightarrow [2]$, r is a functor $p/b \longrightarrow p^{-1}(b)$.

For $(e, w) \in (p/b)_0$ and $e' \in p^{-1}(b)_0$,

$$\varphi : \mathrm{Hom}_{p/b}((e, w), i(e')) \longrightarrow \mathrm{Hom}_{p^{-1}(b)}(r(e, w), e')$$

is given as follow. Since $i(e') = (e', 1_b)$, then an element $f \in \mathrm{Hom}_{p/b}((e, w), i(e'))$ is a morphism $f : e \longrightarrow e'$ such that $p(f) = w$. Then, there exists a morphism $g : \tilde{w}(1) \longrightarrow e'$ such that $p(g) = 1_b$ and $g \circ \tilde{w} = f$ since the lift property of p for J_2. Define φ as $\varphi(f) = g$. Conversely,

$$\chi : \mathrm{Hom}_{p^{-1}(b)}(r(e, w), e') \longrightarrow \mathrm{Hom}_{p/b}((e, w), i(e'))$$

is defined by $h \mapsto h \circ \tilde{w}$. It is obvious that $\chi \circ \varphi = 1$. And $\varphi \circ \chi = 1$ since the lift property of p for J_3. Thus p is a precofibered category. Similarly, we can prove that p is a prefibered category. ☐

Definition 5.8. Let $f : X \longrightarrow Y$ be a prefibered category, then define a functor $\alpha^* : f^{-1}(y') \longrightarrow f^{-1}(y)$ for a morphism $\alpha : y \longrightarrow y'$ in Y as follows. α induces a functor $\alpha' : y'/f \longrightarrow y/f$ as $(x, w) \longrightarrow (x, w \circ \alpha)$. Then $\alpha^* = r \circ \alpha' \circ i$, where and $r : y/f \longrightarrow p^{-1}(y)$ is a right adjoint functor of i. Similarly, let $f : X \longrightarrow Y$ be a precofibered category, then define a functor $\alpha_* : f^{-1}(y) \longrightarrow f^{-1}(y')$ for a morphism $\alpha : y \longrightarrow y'$ in Y as follows. α induces a functor $\alpha' : f/y \longrightarrow f/y'$ as $(x, w) \longrightarrow (x, \alpha \circ w)$. Then $\alpha_* = r \circ \alpha' \circ j$, where and $r : f/y' \longrightarrow p^{-1}(y')$ is a left adjoint functor of j.

Proposition 5.9. *Let $p : E \longrightarrow B$ a fibration, then $f_* : p^{-1}(b) \longrightarrow p^{-1}(b')$ and $f^* : p^{-1}(b') \longrightarrow p^{-1}(b)$ satisfy $f_* \circ f^* \cong 1$ and $f^* \circ f_* \cong 1$ for $f : b \longrightarrow b'$ in B.*

Proof. Now, $f_*(e) = \tilde{f}(1)$ where $\tilde{f}$ is a morphism in E such that $p \circ \tilde{f} = f$ and $\tilde{f}(0) = e$. And $f^*(e') = \overline{f}(0)$ where $\overline{f}$ is a morphism in E such that $p \circ \overline{f} = f$ and $\overline{f}(1) = e'$. Natural isomorphisms $f_* \circ f^* \cong 1$ and $f^* \circ f_* \cong 1$ can be defined from the lifting property of p for J_2 and J_3. ☐

Corollary 5.10. *Let $p : E \longrightarrow B$ a covering, then $f_* : p^{-1}(b) \longrightarrow p^{-1}(b')$ is an isomorphism for $f : b \longrightarrow b'$ in B.*

Proof. By Proposition 5.9. ☐

Theorem 5.11. [Qui68] *Suppose $f : X \longrightarrow Y$ is prefibered (resp. precofibered) and that for every arrow $\alpha : y \longrightarrow y'$, $B(\alpha^*) : B\left(f^{-1}(y')\right) \longrightarrow B\left(f^{-1}(y)\right)$ (resp. $B(\alpha_*) : B\left(f^{-1}(y)\right) \longrightarrow B\left(f^{-1}(y')\right)$) is a homotopy equivalence. Then for any $x \in f^{-1}(y)$, we have an exact sequence of homotopy groups*

$$\cdots \longrightarrow \pi_{i+1}(BY, y) \longrightarrow \pi_i\left(B\left(f^{-1}(y)\right), x\right) \longrightarrow \pi_i(BX, x) \xrightarrow{f_*} \pi_i(BY, y) \longrightarrow \cdots$$

Corollary 5.12. *Let $p : X \longrightarrow Y$ be a fibration in* **Cat**$_1$*, then for any $x \in f^{-1}(y)$, we have an exact sequence of homotopy groups*

$$\cdots \longrightarrow \pi_{i+1}(BY, y) \longrightarrow \pi_i\left(B\left(f^{-1}(y)\right), x\right) \longrightarrow \pi_i(BX, x) \xrightarrow{f_*} \pi_i(BY, y) \longrightarrow \cdots$$

Proof. This follows from Proposition 5.7 and 5.9 and Theorem 5.11. ☐

Theorem 5.13. *p is a fibration in* **Cat**$_1$ *iff Np is a Kan fibration.*

Proof. Let $p : E \longrightarrow B$ be a fibration, we will prove that Np is a Kan fibration. Given a following diagram in **SSets**

$$\begin{tikzcd}
\Lambda^n_k \arrow[r] \arrow[d] & NE \arrow[d, "Np"] \\
\Delta[n] \arrow[r] & NB
\end{tikzcd}$$

for $0 < n$ and $0 \leqq k \leqq n$. Since (c, N) is an adjoint pair, the above diagram induce a next commutative diagram in **Cat**.

$$\begin{tikzcd}
c\Lambda^n_k \arrow[r] \arrow[d] & E \arrow[d, "p"] \\
c\Delta[n] \cong [n] \arrow[r] & B
\end{tikzcd}$$

Now the left vertical arrow is $c(\Lambda^n_k) \cong [n]$ for $n \geqq 4$, and for $0 < n \leqq 3$, this is pushout of a morphism of generating trivial cofibration J in the 1-type model structure. Then there exists a morphism $h : [n] \longrightarrow E$ such that commuting following diagram.

$$\begin{tikzcd}
c\Lambda^n_k \arrow[r] \arrow[d] & E \arrow[d, "p"] \\
c\Delta[n] \cong [n] \arrow[r] \arrow[ur, "h"] & B
\end{tikzcd}$$

Thus, $h^\sharp$ the adjoint morphism of h makes the following diagram commutative.

$$\begin{tikzcd}
\Lambda^n_k \arrow[r] \arrow[d] & NE \arrow[d, "Np"] \\
\Delta[n] \arrow[r] \arrow[ur, "h^\sharp"] & NB
\end{tikzcd}$$

Therefore, Np is a Kan fibration.

Conversely, let Np be a Kan fibration. Given a following diagram in **Cat**,

$$\begin{tikzcd}
X \arrow[r] \arrow[d] & E \arrow[d, "p"] \\
Y \arrow[r] & B
\end{tikzcd}$$

where, $X \longrightarrow Y$ belongs in J. Since $cN \cong 1_{\mathbf{Cat}}$, it suffices to show that there exists a morphism $NY \longrightarrow NE$ commuting the following diagram in **SSets** applied N for the above diagram.

$$\begin{tikzcd}
NX \arrow[r] \arrow[d] & NE \arrow[d, "Np"] \\
NY \arrow[r] & NB
\end{tikzcd}$$

When $X \longrightarrow Y$ belongs J_1 or J_3, then $NX \longrightarrow NY$ is $\Lambda^1_{0(1)} \longrightarrow \Delta[1]$ or $\Lambda^2_{0(2)} \longrightarrow \Delta[2]$, respectively. Since Np is a Kan fibration, then there is a morphism $NY \longrightarrow NE$ what we want. Now suppose $X \longrightarrow Y$ is $k : CS^1 \longrightarrow [2]$, then consider a following commutative diagram

$$\begin{tikzcd}
\Lambda^3_0 \arrow[r, "r"] \arrow[d] & N(CS^1) \arrow[r] \arrow[d, "Nk"'] & NE \arrow[d, "Np"] \\
\Delta[3] \arrow[r, "Nd"'] & N([2]) \arrow[r] & NB
\end{tikzcd}$$

where $d : [3] \longrightarrow [2]$ is defined by $m \mapsto m$ for $0 \leqq m \leqq 2$ and $3 \mapsto 2$. And

$$r : \Lambda^3_0 \longrightarrow N(CS^1)$$

is a canonical projection commuting the diagram. d is a retraction of $k : [2] \longrightarrow [3]$ given by $0 \mapsto 0$, $1 \mapsto 1$, $2 \mapsto 3$. Because Np is a Kan fibration, then there exists a morphism $h : \Delta[3] \longrightarrow NE$ such that

$$\begin{tikzcd}
\Lambda^3_0 \arrow[r, "r"] \arrow[d] & N(CS^1) \arrow[r] & NE \arrow[d, "Np"] \\
\Delta[3] \arrow[r, "Nd"'] \arrow[urr, "h"] & N([2]) \arrow[r] & NB
\end{tikzcd}$$

is commutative. Apply the functor c, then we obtain a following diagram.

$$\begin{tikzcd}
c\Lambda^3_0 \arrow[r, "cr"] \arrow[d] & CS^1 \arrow[r] & E \arrow[d, "p"] \\
{[3]} \arrow[r, "d"'] \arrow[urr, "ch"] & {[2]} \arrow[r] & B
\end{tikzcd}$$

Now $c\Lambda^3_0$ is formed as follows,

$$\begin{tikzcd}
0 \arrow[r] & 1 \arrow[rr, bend left] \arrow[dr] & & 3 \\
& & 2 \arrow[ur, "f"'] &
\end{tikzcd}$$

and cr carries f to the identity map. Thus cr is a retraction of the inclusion functor $CS^1 \hookrightarrow c\Lambda^3_0$, then $ch \circ j : [2] \longrightarrow E$ makes the following diagram commutative.

$$\begin{tikzcd}
CS^1 \arrow[r] \arrow[d] & E \arrow[d, "p"] \\
{[2]} \arrow[r] \arrow[ur, "ch\circ j"] & B
\end{tikzcd}$$

Thus p is a fibration. $\square$

Corollary 5.14. *G is a groupoid iff $N(G)$ is a Kan complex.*

Proof. By Proposition 5.2, groupoids coincide with fibrant objects in the category of small categories equipped with 1-type model structure, and Kan complexes coincides with fibrant objects in the category of simplicial sets equipped with Kan type model structure [Hov99]. Thus Theorem 5.13 implies that G is a groupoid iff $N(G)$ is a Kan complex since N preserves terminal objects. $\square$

The functorial factorizations in the 1-type model structure on **Cat** is induced from the small objects argument in [Hir03]. However, the small objects argument is too abstract and difficult. Now, we define another functorial factorizations on $\mathbf{Cat}_1$ which induce the groupoidification in Definition 2.2 and the universal cover in Definition 4.6.

Definition 5.15. For a functor $f : X \longrightarrow Y$, define a category E_f as follows.

$$(E_f)_0 = \{(x, y_*) \in X_0 \times \mathrm{Mor}(\pi(Y))_0 \mid f(x) = y_0\}$$

and

$$\mathrm{Hom}_{E_f}((x, y_*), (x', y'_*)) = \{(g_*, g) \in \mathrm{Hom}_{\pi(X)}(x, x') \times \mathrm{Hom}_Y(y_n, y'_m) \mid y'_* \circ f(g_*) \circ y'_* = g \circ y_* \in \pi(Y)\}$$

where y_n and y'_m are last objects of y_*, y'_*, respectively. When $X = *$, then E_f is $\widehat{Y}$ in Definition 4.6. We define a functor $j : X \longrightarrow E_f$ as $x \mapsto (x, 1_{f(x)})$ and a functor $p : E_f \longrightarrow Y$ as $(x, y_*) \mapsto y_n$ where y_n is the last object of y_*. Define $\alpha, \beta : \mathrm{Mor}(\mathbf{Cat}) \longrightarrow \mathrm{Mor}(\mathbf{Cat})$ as $\alpha(f) = j$ and $\beta(f) = p$, then it is obvious that $f = \beta(f) \circ \alpha(f)$ and (α, β) is a functorial factorization of **Cat**.

Proposition 5.16. *$p : E_f \longrightarrow Y$ in Definition 5.15 is a fibration for any $f : X \longrightarrow Y$.*

Proof. Given a commutative diagram.

$$\begin{tikzcd}
* \arrow[r, "\alpha"] \arrow[d] & E_f \arrow[d, "p"] \\
{[1]} \arrow[r, "\beta"'] & Y
\end{tikzcd}$$

Let $\alpha(*) = (x, y_*)$, then $\beta(0 \longrightarrow 1) : \beta(0) = y_n \longrightarrow \beta(1)$. Thus, we define a functor $\gamma : [1] \longrightarrow E_f$ by $\gamma(0) = \alpha(*) = (x, y_*)$, $\gamma(1) = (x, \beta(0 \longrightarrow 1) \circ y_*)$ and $\gamma(0 \longrightarrow 1) = (1, \beta(0 \longrightarrow 1))$. γ makes the following diagram commutative,

$$\begin{tikzcd}
* \arrow[r, "\alpha"] \arrow[d] & E_f \arrow[d, "p"] \\
{[1]} \arrow[r, "\beta"'] \arrow[ur, "\gamma"] & Y
\end{tikzcd}$$

then p has the lifting property for J_1. Next, given the following commutative diagram.

$$\begin{tikzcd}
I_2 \arrow[r, "\alpha"] \arrow[d] & E_f \arrow[d, "p"] \\
{[2]} \arrow[r, "\beta"'] & Y
\end{tikzcd}$$

Then, the image of α represents a diagram in E_f as

$$(x', y'_*) \overset{(g_*, g)}{\longleftarrow} (x, y_*) \overset{(h_*, h)}{\longrightarrow} (x'', y''_*).$$

Now $\beta(1 \longrightarrow 2)$ is a morphism from last point of y'_* to y''_*. Thus, we obtain a morphism

$$(h_* \circ g_*^{-1}, \beta(1 \longrightarrow 2)) : (x', y'_*) \longrightarrow (x'', y''_*)$$

and it gives a functor $[2] \longrightarrow E_f$ commuting the above diagram, then p has the lifting property for J_2.

At last, given the following commutative diagram.

$$\begin{tikzcd}
CS^1 \arrow[r, "\alpha"] \arrow[d] & E_f \arrow[d, "p"] \\
{[2]} \arrow[r, "\beta"'] & Y
\end{tikzcd}$$

then, the image of α represents a diagram in E_f as

$$\begin{tikzcd}
(x, y_*) \arrow[r, "{(g_*, g)}"] & (x', y'_*) \arrow[r, shift left, "{(h_*, h)}"] \arrow[r, shift right, "{(h'_*, h')}"'] & (x'', y''_*)
\end{tikzcd}$$

Since $h_* \circ g_* = h'_* \circ g_*$ in $\pi(X)$, then

$$h_* = h_* \circ g_* \circ g_*^{-1} = h'_* \circ g_* \circ g_*^{-1} = h'_*$$

in $\pi(X)$. Moreover, β implies that $h = h'$, then $(h_*, h) = (h'_*, h')$ and it gives a functor $[2] \longrightarrow E_f$ commuting the above diagram, then p has the lifting property for J_3. □

Proposition 5.17. *$j : X \longrightarrow E_f$ in Definition 5.15 is a trivial cofibration for any $f : X \longrightarrow Y$.*

Proof. It is obvious that j is a cofibration, so we will show that j is a weak 1-equivalence. First, we see $j_* : \pi_0(X) \longrightarrow \pi_0(E_f)$.

We take an element $[x, y_*] \in \pi_0(E_f)$. (x, y_*) represents the diagram

$$f(x) = y_0 \longrightarrow y_1 \longleftarrow y_2 \longrightarrow \cdots \longleftarrow y_n$$

then, we consider the next commutative diagram.

$$\begin{tikzcd}
f(x) \arrow[r, "="] \arrow[dr] & f(x) \arrow[d] & f(x) \arrow[l, "="'] \arrow[r, "="] \arrow[d] & \cdots \\
 & y_1 & y_1 \arrow[l, "="'] \arrow[r, "="] & \cdots \\
 & & y_2 \arrow[u] \arrow[ul] \arrow[r] \arrow[dr] & \cdots \\
 & & & \cdots
\end{tikzcd}$$

When we regard vertical sequences as objects in E_f, the above diagram implies that

$$j_*[x] = [x, 1_{f(x)}] = [x, y_*]$$

then j_* is surjective. On the other hand, for $[x], [y] \in \pi_0(X)$, assume that $j_*[x] = j_*[y] \in \pi_0(E_f)$. It is obvious that $[x] = [y] \in \pi_0(X)$ from the definition of set of morphisms of E_f. Then j_* is injective.

Next we see the induced map $j_* : \pi_1(X, x) \longrightarrow \pi_1(X, (x, 1_{f(x)}))$. We take an element

$$\sigma : (x, 1_{f(x)}) \xrightarrow{(g_{1,*},g_1)} (x_1, y_{1,*}) \xleftarrow{(g_{1,*},g_1)} (x_2, y_{2,*}) \longrightarrow \cdots \xleftarrow{(g_{n,*},g_n)} (x_n, y_{n,*}) = (x, 1_{f(x)})$$

of $\pi_1(X, (x, 1_{f(x)}))$. For the i-th morphism $(g_{i,*}, g_i) : (x_{i-1}, y_{i-1,*}) \longrightarrow (x_i, y_{i,*})$ (in the case of i even), $g_{i,*}$ is an element of $\mathrm{Hom}_{\pi(X)}(x_{i-1}, x_i)$. Let

$$\rho = g_{n,*}^{-1} \circ \cdots \circ g_{2,*}^{-1} \circ g_{1,*} \in \pi_1(X, x),$$

then $j_*(\rho) = \sigma$ in $\pi_1(X)$. Therefore j_* is surjective. At last,

$$\mathrm{Ker}\ (j_*) = \{x_* \in \pi_1(X, x) \mid j_*(x_*) = 0 \in \pi_1(E_f, (x, 1_{f(x)})\} = 0$$

since the both morphisms and compositions in E_f are same as X and the definition of j. Thus j_* is injective. □

Corollary 5.18. *The functorial factorization (α, β) in Definition 5.15 satisfies the axiom 4 of the 1-type model structure on* **Cat**.

Proof. By Proposition 5.16 and Proposition 5.17. □

Remark 5.19. *Let C be a category, and denote $q : C \longrightarrow *$ the morphism from C to the terminal object $*$ in* **Cat**. *Then $\alpha(q)$ is the functor $C \longrightarrow \pi(C)$ in Definition 2.2, where (α, β) is the functorial factorization in Definition 5.15.*

Remark 5.20. *Let $(C, *)$ be a pointed and path connected category, and denote $k : * \longrightarrow C$ as the inclusion functor. Then $\beta(k)$ is the universal cover $\widehat{C} \longrightarrow C$ in Definition 4.6, where (α, β) is the functorial factorization in Definition 5.15.*

6 The classification theorem of coverings of small categories

We show that a subgroup H of $\pi_1(C)$ corresponds to a covering over C with the fundamental group of the total category isomorphic to H. This is a generalization of the classification of coverings in the category of groupoids in [May99]

Definition 6.1. Let G be a group and C be a small category. C is called a G-category if G acts on C_0 and there is a correspondence of the set of morphism $\mathrm{Hom}_C(x, y) \longrightarrow \mathrm{Hom}_C(gx, gy)$ for $g \in G$ satisfying the associativity and unit conditions and composition identity. This is regarded as a functor $G \longrightarrow$ **Cat** such that $* \mapsto C$, where $*$ is an only object of G. In particular, C is called a free G-category when the action of G on C_0 is free.

Definition 6.2. [CM06] Let G be a group and C be a free G-category. Define C/G as $(G/C)_0 = C_0/G$ and

$$\mathrm{Hom}_{C/G}(\alpha, \beta) = \coprod_{x \in \alpha, y \in \beta} \mathrm{Hom}_C(x, y)/G$$

where G-action is given by $(g, f) \mapsto g(f)$. The composition

$$\mathrm{Hom}_{C/G}(\alpha, \beta) \times \mathrm{Hom}_{C/G}(\beta, \gamma) \longrightarrow \mathrm{Hom}_{C/G}(\alpha, \gamma)$$

is defined as $([\varphi], [\chi]) \mapsto [g(\chi) \circ \varphi]$ where $g \in G$ such that $t(\varphi) = gs(\chi)$. Because of the freeness of the action, C/G is a small category.

Remark 6.3. *If a free G-category C is regarded as a functor $G \longrightarrow$ **Cat**, then the colimit of the functor coincides with C/G.*

Lemma 6.4. *Let C be a free G-category, then the canonical projection $p : C \longrightarrow C/G$ is a covering.*

Proof. Given a commutative diagram

$$\begin{tikzcd}
* \arrow[r, "c"] \arrow[d] & C \arrow[d, "p"] \\
{[1]} \arrow[r, "g"'] & C/G
\end{tikzcd}$$

Now $g(0 \longrightarrow 1) : g(0) = [c] \longrightarrow g(1)$ and

$$\mathrm{Hom}_{C/G}([c], g(1)) = \coprod_{x \in [c], y \in g(1)} \mathrm{Hom}_C(x, y)/G,$$

then $g(0 \longrightarrow 1) = [f]$ where $f \in \mathrm{Hom}_C(x, y)$. Since $x \in [c]$, there exists an element $g \in G$ such that $gx = c$, so $g(f) \in \mathrm{Hom}_C(gx = c, gy)$. We define a functor $h : [1] \longrightarrow C$ by $h(0) = c, h(1) = gy, h(0 \longrightarrow 1) = g(f)$.

$$p \circ h(0 \longrightarrow 1) = [h(0 \longrightarrow 1)] = [g(f)] = [f] = g(0 \longrightarrow 1).$$

On the other hand, suppose there exists a morphism $h' : [1] \longrightarrow C$ satisfying the same property as h. Since $h'(0) = c$, then $h'(0 \longrightarrow 1) \in \mathrm{Hom}(c, z)$. Because of $[h(0 \longrightarrow 1)] = [h'(0 \longrightarrow 1)]$, there exists an element $g \in G$ such that $g(h(0 \longrightarrow 1)) = h'(0 \longrightarrow 1)$. By looking at the source object, the above equation implies $gc = c$. Since the G-action is free, we obtain $g = e$. Thus $h(0 \longrightarrow 1) = h'(0 \longrightarrow 1)$ and $h = h'$. Similarly, p has the unique lifting property for $1 : 0 \longrightarrow [1]$. ☐

Definition 6.5. Let $(C, *)$ be a pointed small category. Define a $\pi_1(C)$-action on $\widehat{C}$

$$\mu : \pi_1(C) \times \widehat{C} \longrightarrow \widehat{C}$$

by concatenating the sequences on object and inclusion on the set of morphisms.

Lemma 6.6. *The action $\mu : \pi_1(C) \times \widehat{C} \longrightarrow \widehat{C}$ in Definition 6.5 is free.*

Proof. For $x_* = (* = x_0 \longrightarrow x_1 \longleftarrow x_2 \longrightarrow \cdots \longleftarrow x_n = *) \in \pi_1(C)$, assume $\mu(y_*, x_*) = x_*$, then $y_* \circ x_* = x_*$ and $y_* = x_* \circ x_*^{-1} = 1$ in $\pi_1(C)$. ☐

Definition 6.7. Let $(C, *)$ be a pointed small category and H be any subgroup of $\pi_1(C)$. Since lemma 6.6, H acts free on $\widehat{C}$ as restricted action of Definition 6.5. $T : \widehat{C} \longrightarrow C$ induces a morphism $\widetilde{T} : \widehat{C}/H \longrightarrow C$ making the following diagram commutative.

$$\begin{tikzcd}
\widehat{C} \arrow[rr, "p"] \arrow[dr, "T"'] & & \widehat{C}/H \arrow[dl, "\widetilde{T}"] \\
& C &
\end{tikzcd}$$

Lemma 6.8. *$\widetilde{T} : \widehat{C}/H \longrightarrow C$ in Definition 6.7 is a covering for any subgroup H of $\pi_1(C)$.*

Proof. The action of H is to circle around the starting point of a sequence. On the other hand, T is a correspondence which assigns the last point of a sequence. Then the H-action does not have an influence on T, so we can show $\widetilde{T}$ is a covering by a similar argument we used to proof of Proposition 4.7. $\square$

Proposition 6.9. $\widetilde{T}_*(\pi_1(\widehat{C}/H)) = H$

Proof. For $h_* = (* = h_0 \longrightarrow h_1 \longleftarrow h_2 \longrightarrow \cdots \longleftarrow h_n = *) \in H \subset \pi_1(C)$ and $0 \leqq m \leqq n$, define $h_{*,m}$ as a subsequence of h_* with length m, then $h_{*,m} \in \widehat{C}$ for all m. Then we obtain an element

$$[h_*]_* = [h_{*,*}] \in \pi_1(\widehat{C}/H)$$

and $\widetilde{T}'_*[h_*]_*) = h_*$. Thus $H \subset \widetilde{T}_*(\pi_1(\widehat{C}/H))$.

Conversely, for $x_* \in \widetilde{T}_*\pi_1(\widehat{C}/H)$, there exists an element $[c_*]_* \in \pi_1(\widehat{C}/H)$ such that $\widetilde{T}_*[c_*]_* = x_*$. Take each of the last objects of

$$\begin{tikzcd}
{[c_*]_0} \arrow[r, "{[f_1]}"] & {[c_*]_1} & {[c_*]_2} \arrow[l, "{[f_2]}"'] \arrow[r, "{[f_3]}"] & {[c_*]_3} & \cdots \arrow[l] & {[c_*]_n} \arrow[l]
\end{tikzcd}$$

then,

$$(c_{m_0} \xrightarrow{f_1} c_{m_1} \xleftarrow{f_2} c_{m_3} \xrightarrow{f_3} \cdots \longleftarrow c_{m_n}) = (x_0 \longrightarrow x_1 \longleftarrow x_2 \longrightarrow \cdots \longleftarrow x_l)$$

By the definition of the morphism of $\pi(\widehat{C}/H)$, $[f_{i+1}] \circ [c_*]_i = [c_*]_{i+1}$ in $\pi(\widehat{C}/H)$. Then there exists an element $h_{*,i} \in H$ such that $f_{i+1} \circ c_{*,i} = c_{*,i+1} \circ h_{*,i}$ in $\pi(\widehat{C})$. Using a similarly method of proposition 4.8, we obtain

$$x_* = (c_{m_0} \xrightarrow{f_1} c_{m_1} \xleftarrow{f_2} c_{m_3} \xrightarrow{f_3} \cdots \longleftarrow c_{m_n}) = h_{*,1}h_{*,2}^{-1} \cdots h_{*,n}^{(-1)^{n-1}} \in H$$

thus $\widetilde{T}_*\pi_1(\widehat{C}/H) \subset H$. $\square$

Definition 6.10. Let $(B, *)$ be a pointed small category, define the category $\mathrm{Cov}_*(B)$ of coverings with base point as follows.

$$\mathrm{Cov}_*(B)_0 = \{(p, e) \mid p \text{ is a covering over } B\ ,\ e \in p^{-1}(*)\}.$$

A morphism between $(p : E \longrightarrow B; e)$ and $(q : E' \longrightarrow B; e')$ is a functor $f : E \longrightarrow E'$ such that $f(e) = e'$ and $q \circ f = p$. The set of isomorphic classes of the set of objects of $\mathrm{Cov}_*(B)$ is denoted by $\overline{\mathrm{Cov}}_*(B)$.

On the other hand, when G is a group, we denote the set of subgroups of G by $O(G)$.

Lemma 6.11. *Let $p : E \longrightarrow B$ be a covering and $f : X \longrightarrow B$ be a functor and assume X is path connected. Choose $b_0 \in B_0, e_0 \in p^{-1}(b_0), x_0 \in f^{-1}(b_0)$, then there exists a unique functor $g : X \longrightarrow E$ making the following diagram commutative*

$$\begin{tikzcd}
X \arrow[rr, dotted, "g"] \arrow[dr, "f"'] & & E \arrow[dl, "p"] \\
& B &
\end{tikzcd}$$

and $g(x_0) = e_0$ iff $f_*(\pi_1(X, x_0)) \subset p_*(\pi_1(E, e_0))$ in $\pi_1(B, b_0)$.

Proof. If there exists a functor $g : X \longrightarrow E$ such that $p \circ g = f$ and $g(x_0) = e_0$, then

$$f_*(\pi_1(X, x_0)) = p_*(g_*(\pi_1(X, x_0))) \subset p_*(\pi_1(E, e_0)).$$

Conversely, assume that $f_*(\pi_1(X, x_0)) \subset p_*(\pi_1(E, e_0))$. Take an object $x \in X_0$, then there exists a zigzag sequence x_* from x_0 to x since X is path connected. Since p is a covering, there exists a unique morphism $e_* : I_n \longrightarrow E$ making the following diagram commutative.

$$\begin{tikzcd}
* \arrow[r, "e_0"] \arrow[d] & E \arrow[d, "p"] \\
I_n \arrow[r, "f(x_*)"'] \arrow[ur, "e_*"] & B
\end{tikzcd}$$

Define $g : X \longrightarrow E$ by $g(x) = T(e_*)$ on object. $g(x)$ does not depend on the choice of x_* from [May99].

Next for $\alpha : x \longrightarrow y$, assume that x_* is a zigzag sequence between x_0 and x. We can take a zigzag sequence between x_0 and y as $\alpha \circ x_*$.

Then we get a morphism $g(x) \longrightarrow g(y)$ of E over $f(\alpha)$ from the lifting property. It gives a correspondence of g on the set of morphisms. The universality of g is shown from the unique lifting property of p. □

Corollary 6.12. *Let $(p : E \longrightarrow B; e)$ and $(p' : E' \longrightarrow B; e')$ be elements of $Cov_*(B)$ If*

$$p_*(\pi_1(E, e)) = p'_*(\pi_1(E', e'))$$

in $\pi_1(B, *)$ then (p, e) is isomorphic to (p', e').

Proof. By the universality of existence of a functor $E \longrightarrow E'$ from lemma 6.11. □

Definition 6.13. Let $(C, *)$ be a pointed category, define a map

$$F : O(\pi_1(C, *)) \longrightarrow \mathrm{Cov}_*(C)_0$$

by $H \mapsto (\widehat{C}/H \longrightarrow C; *)$. And define a map

$$G : \mathrm{Cov}_*(C)_0 \longrightarrow O(\pi_1(C, *))$$

by $(p : E \longrightarrow C; e) \mapsto p_*(\pi_1(E, e))$.

Lemma 6.14. *If $(p : E \longrightarrow C; e) \cong (p' : E' \longrightarrow C; e')$, then $G(p) = G(p')$.*

Proof. By assumption, there exists an isomorphism $f : E \longrightarrow E'$ such that $f(e) = e'$ and

$$\begin{tikzcd}
X \arrow[rr, "f"] \arrow[dr, "p"'] & & E \arrow[dl, "p'"] \\
& B &
\end{tikzcd}$$

is a commutative, then $G(p) = p_*(\pi_1(E, e)) = p'_*(f_*(\pi_1(E, e)) = p'_*(\pi_1(E', f(e))) = G(p')$. □

Definition 6.15. Define a map

$$\widetilde{F} : O(\pi_1(C, *)) \xrightarrow{F} \mathrm{Cov}_*(C)_0 \xrightarrow{p} \overline{\mathrm{Cov}}_*(C)$$

where p is the projection and F is the map in Definition 6.13. And

$$\widetilde{G} : \overline{\mathrm{Cov}}_*(C) \longrightarrow O(\pi_1(C, *)).$$

is a map induced from G in Definition 6.13 by Lemma 6.14.

Theorem 6.16. *$\widetilde{G}$ is an inverse map of $\widetilde{F}$.*

Proof. By Proposition 6.9, it is clearly that $\widetilde{G} \circ \widetilde{F} = 1$. We will show that $\widetilde{F} \circ \widetilde{G} = 1$. Let $p : E \longrightarrow C$ be a covering and $e \in p^{-1}(*)$. Consider the covering $\widetilde{T} : \widehat{C}/\left(p_*\pi_1(E, e)\right) \longrightarrow C$ then Proposition 6.9 suggests that

$$p_*(\pi_1(E, e)) = \widetilde{T}_*\left(\pi_1\left(\widehat{C}/\left(p_*(\pi_1(E, e))\right)\right)\right).$$

Then Corollary 6.12 implies that (p, e) and $(\widetilde{T}, *)$ are isomorphic, thus $\widetilde{F} \circ \widetilde{G} = 1$. □

7 Relations between the 1-type model category and other model categories

Cat with the 1-type model structure is related to other model categories by the following pairs of adjoint functors.

$$\begin{tikzcd}
\mathbf{Grpd} \arrow[r, shift right, "i"'] & \mathbf{Cat} \arrow[l, shift right, "\pi"'] \arrow[r, shift right, "N"'] & \mathbf{SSets} \arrow[l, shift right, "c"'] \arrow[r, shift right, "|-|"'] & \mathbf{Spaces} \arrow[l, shift right, "S"']
\end{tikzcd}$$

Definition 7.1. [Hir03] Let M and N be model categories and let

$$F : M \Longleftrightarrow N : G$$

be a pair of adjoint functors. We say that (F, G) is a Quillen pair if F preserves cofibrations and G preserves fibrations. Furthermore (F, G) is called a pair of Quillen equivalences if for every cofibrant object X in M, every fibrant object Y in N, and every map $f : X \longrightarrow GY$ in M, the map f is a weak equivalence in M iff the corresponding map $f^\sharp : FX \longrightarrow Y$ is a weak equivalence in N.

Proposition 7.2. [Hir03] *Let M and N be model categories and let $F : M \Longleftrightarrow N : G$ be a pair of adjoint functors. Then the following are equivalent:*

1. *(F, G) is a Quillen pair.*
2. *F preserves both cofibrations and trivial cofibrations.*
3. *G preserves both fibrations and trivial fibrations.*

Proposition 7.3.

$$\pi : \mathbf{Cat}_1 \Longleftrightarrow \mathbf{Grpd} : i$$

is a pair of Quillen equivalences, where **Grpd** *is the category of groupoids with the Anderson-type model structure in Theorem 3.9.*

Proof. Because π preserves weak equivalences and cofibrations, Proposition 7.2 implies that (π, i) is a Quillen pair, furthermore, (π, i) is a pair of Quillen equivalences since the inclusion functor $C \longrightarrow \pi(C)$ in Definition 2.2 is a weak 1-equivalence for any small category C. ☐

Proposition 7.4. *Denote a category of small categories with the Joyal and Tierney-type model category by $\mathbf{Cat}_{JT}$ in Theorem 3.8. Then*

$$1 : \mathbf{Cat}_{JT} \Longleftrightarrow \mathbf{Cat}_1 : 1$$

is a Quillen pair.

Proof. Because a cofibrations and a weak equivalence in $\mathbf{Cat}_{JT}$ is a cofibration and a weak equivalence in $\mathbf{Cat}_1$, respectively. Proposition 7.2 implies that $(1, 1)$ is a Quillen pair. ☐

Proposition 7.5. *Denote the category of simplicial sets equipped with the Kan type model structure by $\mathbf{SSets}_K$ in [Hov99], then*

$$c : \mathbf{SSets}_K \Longleftrightarrow \mathbf{Cat}_1 : N$$

is a Quillen pair.

Proof. Theorem 5.13 implies that N preserves fibrations. Because a cofibration $i : A \longrightarrow B$ in $\mathbf{SSets}_K$ is injective for all dimensions, in particular, $i_0 : A_0 \longrightarrow B_0$ is injective. Then the map on the set of objects of $ci : cA \longrightarrow cB$ coincides with $i_0 : A_0 \longrightarrow B_0$, thus ci is a cofibration. Therefore c preserves cofibrations, then (c, N) is a Quillen pair. ☐

We will prove that $\mathbf{Cat}_1$ is Quillen equivalent to the category of simplicial set with the 1-type model category in Theorem 3.19. From Definition 3.14, a weak 1-equivalence $f : X \longrightarrow Y$ in **SSets** is a map inducing isomorphisms

$$|f|_* : \pi_n(|X|, x) \longrightarrow \pi_n(|Y|, |f|(x))$$

for $n = 0, 1$ and $x \in |X|$. In [May92], the fundamental (homotopy) group of a fibrant simplicial set X in $\mathbf{SSets}_K$ is defined, and proved that $\pi_1(X, *) \cong \pi_1(|X|, *)$ in [May92]. Those fibrant simplicial sets are called Kan complexes.

Definition 7.6. Let X be a simplicial set. Two 1-simplices $x, y \in X_1$ is homotopic, denoted $x \simeq y$, if $d_i x = d_i y$ for $i = 0, 1$, and there exists a 2-simplex $z \in X_2$ such that $d_0 z = s_0 d_0 x$, $d_1 z = x, d_2 z = y$. z is called a homotopy from x to y. $\simeq$ is an equivalence relation on X_1, when X is a Kan complex. Let X be a Kan complex and $*$ be a vertex of X, define $\pi_1(X, *) = \{x \in X_1 \mid d_0 x = d_1 x = *\}/\simeq$.

Lemma 7.7. [May92] *There exist a group structure on $\pi_1(X, *)$ and*

$$\theta_* : \pi_1(X.*) \longrightarrow \pi_1(S|X|, *) = \pi_1(|X|, *)$$

is an isomorphism of groups, where $\theta : X \longrightarrow S|X|$ is the counit map of the adjoint pair $(|-|, S)$.

Lemma 7.8. *Let X be a Kan complex and $*$ be a vertex of X, then*

$$\eta_* : \pi_1(|X|, *) \longrightarrow \pi_1(|NcX|, *)$$

is an isomorphism, where $\eta : X \longrightarrow NcX$ is the counit map of the adjoint pair (c, N).

Proof. Define a map

$$\varepsilon : X \xrightarrow{\eta} NcX \xrightarrow{Ni} N\pi cX.$$

where $i : cX \longrightarrow \pi cX$ is the functor in Definition 2.2. Because i is a weak 1-equivalence, it suffices to prove that

$$\varepsilon_* : \pi_1(|X|, *) \longrightarrow \pi_1(|N\pi cX|, *)$$

is an isomorphism. πcX is a groupoid, then $N\pi cX$ is a Kan complex by Corollary 5.2. By lemma 7.7, we will show that

$$\varepsilon_* : \pi_1(X, *) \longrightarrow \pi_1(N\pi cX, *)$$

is an isomorphism. For $[e] \in \pi_1(N\pi cX, *)$, e is a 1-simplex of $N\pi cX$, then e is a morphism of πcX. From the definition of π in Definition 2.2, e is represented as a zigzag sequence of cX as follows.

$$* \xrightarrow{e_1} \cdot \xleftarrow{e_2} \cdot \xrightarrow{e_3} \cdots \xrightarrow{e_n} *$$

e_i is a morphism of cX, then e_i is represented by

$$e_i = e_{i1} e_{i2} \cdots e_{im_i}$$

where $e_{ij} \in X_1$. Since X is a Kan complex, then there exists a $e_i' \in X_1$ such that $e_i = e_i'$ in cX for $0 \leqq i \leqq n$. Then e is

$$* \xrightarrow{e_1'} \cdot \xleftarrow{e_2'} \cdot \xrightarrow{e_3'} \cdots \xrightarrow{e_n'} *.$$

Now X is a Kan complex, there exists an element $d_1 \in X_1$ such that $e_2' \circ d_1 = e_1'$in cX. Next there exists an element $d_2 \in X_1$ such that $e_3' \circ d_1 = d_2$. Repeating this operation, we obtain $d_{n-1} \in X_1$ such that $e_n' \circ d_{n-2} = d_{n-1}$ and $[d_{n-1}] \in \pi_1(X, *)$ and $\varepsilon_*[d_{n-1}] = [e]$.

$$\begin{tikzcd}
* \arrow[r, "e_1'"] \arrow[rr, dotted, bend right, "d_1"'] \arrow[rrr, dotted, bend right=50, "d_2"'] \arrow[rrrr, dotted, bend right=60, "d_{n-1}"'] & \cdot & \cdot \arrow[l, "e_2'"'] \arrow[r, "e_3'"] & \cdots \arrow[r, "e_n'"] & *
\end{tikzcd}$$

Thus ε_* is surjective.

Assume that $\varepsilon_*[e] = 0$ in $\pi_1(Nc\pi X, *)$ for $[e] \in \pi_1(X, *)$. Thus, there exists a 2-simplex (f, g) of $N\pi cX$ such that $d_0(f, g) = 1_*$, $d_1(f, g) = e$, $d_2(f, g) = 1_*$. Then $g = 1_*$, $f = 1_*$, $g \circ f = e$, thus $e = 1_*$ in πcX. There exists the following commutative diagram in cX.

$$\begin{tikzcd}
\ast \arrow[r, "e"] \arrow[d, dash, "="'] & \ast \arrow[d] & \ast \arrow[l, "="'] \arrow[r, "="] \arrow[d] & \cdots & \ast \arrow[l, "="'] \arrow[r, "="] \arrow[d] & \ast \arrow[d, dash, "="] \\
\ast \arrow[r] & \cdot & \cdot \arrow[l] \arrow[r] & \cdots & \cdot \arrow[l] \arrow[r] & \ast \\
\ast \arrow[r] \arrow[u, dash, "="] \arrow[d, dotted, dash] & \cdot \arrow[u] \arrow[d, dotted, dash] & \cdot \arrow[l] \arrow[r] \arrow[u] \arrow[d, dotted, dash] & \cdots & \cdot \arrow[l] \arrow[r] \arrow[u] \arrow[d, dotted, dash] & \ast \arrow[u, dash, "="'] \arrow[d, dotted, dash] \\
\ast \arrow[r] & \cdot & \cdot \arrow[l] \arrow[r] & \cdots & \cdot \arrow[l] \arrow[r] & \ast \\
\ast \arrow[r, "="'] \arrow[u, dash, "="] & \ast \arrow[u] & \ast \arrow[l, "="] \arrow[r] \arrow[u] & \cdots & \cdot \arrow[l] \arrow[r, "="'] \arrow[u] & \ast \arrow[u, dash, "="']
\end{tikzcd}$$

Since X is a Kan complex, we may regard the above morphisms as the 1-simplices of X. Moreover, there exist 1-simplices of X such that the following diagram is commutative.

$$\begin{tikzcd}
\ast \arrow[r, "e"] \arrow[d, dash, "="'] \arrow[rd, dotted] \arrow[rr, dotted, bend left] & \ast \arrow[d] & \ast \arrow[l, "="'] \arrow[r, "="] \arrow[d] \arrow[ld, dotted] \arrow[rd, dotted] \arrow[r, dotted, bend left] & \cdots \arrow[r, dotted, bend left] & \ast \arrow[l, "="'] \arrow[r, "="] \arrow[d] \arrow[ld, dotted] \arrow[rd, dotted] & \ast \arrow[d, dash, "="] \\
\ast \arrow[r] & \cdot & \cdot \arrow[l] \arrow[r] & \cdots \arrow[ll, dotted, bend right] \arrow[rr, dotted, bend left] & \cdot \arrow[l] \arrow[r] & \ast \\
\ast \arrow[r] \arrow[u, dash, "="] \arrow[ru, dotted] \arrow[d, dotted, dash] \arrow[rr, dotted, bend right] & \cdot \arrow[u] \arrow[d, dotted, dash] & \cdot \arrow[l] \arrow[r] \arrow[u] \arrow[lu, dotted] \arrow[ru, dotted] \arrow[d, dotted, dash] \arrow[r, dotted, bend right] & \cdots \arrow[r, dotted, bend right] & \cdot \arrow[l] \arrow[r] \arrow[u] \arrow[lu, dotted] \arrow[ru, dotted] \arrow[d, dotted, dash] & \ast \arrow[u, dash, "="'] \arrow[d, dotted, dash] \\
\ast \arrow[r] & \cdot \arrow[rr, dotted, bend left] & \cdot \arrow[l] \arrow[r] & \cdots \arrow[rr, dotted, bend left] & \cdot \arrow[l] \arrow[r] & \ast \\
\ast \arrow[r, "="'] \arrow[u, dash, "="] \arrow[ru, dotted] \arrow[rr, dotted, bend right] & \ast \arrow[u] & \ast \arrow[l, "="] \arrow[r] \arrow[u] \arrow[lu, dotted] \arrow[ru, dotted] \arrow[r, dotted, bend right] & \cdots \arrow[r, dotted, bend right] & \cdot \arrow[l] \arrow[r, "="'] \arrow[u] \arrow[lu, dotted] \arrow[ru, dotted] & \ast \arrow[u, dash, "="']
\end{tikzcd}$$

It implies that the top of horizontal sequence is the adjacent horizontal sequence in cX. Thus $e = 1_*$ in cX. Therefore, there exists a 2-simplex σ of X such that $d_0\sigma = s_0*$, $d_1\sigma = s_0*$, and $d_2\sigma = e$. Then $[e] = 0$ in $\pi_1(X, *)$, thus ε_* is injective. $\square$

Corollary 7.9. *$\eta : X \longrightarrow NcX$ is a weak 1-equivalence in* **SSets**, *if X is a Kan complex.*

Proof. It is obvious that $\eta_* : \pi_0(X) \longrightarrow \pi_0(NcX)$ is identity map because $X_0 = (NcX)_0$ and the definition of cX. $\eta_* : \pi_1(X, *) \longrightarrow \pi_1(NcX, *)$ is an isomorphism from Lemma 7.8, then η is a weak 1-equivalence. $\square$

Corollary 7.10. *$\eta : X \longrightarrow NcX$ is a weak 1-equivalence in* **SSets** *for all simplicial set X.*

Proof. From the Kan-type model structure on **SSets** in Theorem 3.11, there exists a Kan complex RX and a trivial cofibration $i : X \longrightarrow RX$ in $\mathbf{SSets}_K$ for X. The following diagram

$$\begin{tikzcd}
X \arrow[r, "\eta"] \arrow[d, "i"'] & NcX \arrow[d, "Nci"] \\
RX \arrow[r, "\eta"'] & NcRX
\end{tikzcd}$$

is commutative, and $\eta : RX \longrightarrow NcRX$ is a weak 1-equivalence by Corollary 7.9 and also i is a weak 1-equivalence. By Proposition 7.5,

$$c : \mathbf{SSets}_K \Longleftrightarrow \mathbf{Cat}_1 : N$$

is a Quillen pair, therefore c preserves trivial cofibrations by Proposition 7.2. Then ci is a trivial cofibration in $\mathbf{Cat}_1$, in particular, ci is a weak 1-equivalence in $\mathbf{Cat}$. Therefore Nci is a weak 1-equivalence in $\mathbf{SSets}$, and $\eta : X \longrightarrow NcX$ is a weak 1-equivalence. ☐

Theorem 7.11. *Denote the category of simplicial sets with the* 1*-type model structure in Theorem 3.19 by* $\mathbf{SSets}_1$*, then*

$$c : \mathbf{SSets}_1 \Longleftrightarrow \mathbf{Cat}_1 : N$$

is a pair of Quillen equivalences.

Proof. From Theorem 3.20, (c, N) is a Quillen pair. Suppose X is a cofibrant object in $\mathbf{SSets}_1$ and G is a fibrant object in $\mathbf{Cat}_1$ and $f : X \longrightarrow NG$ is a weak equivalence in $\mathbf{SSets}_1$. $f^\sharp : cX \longrightarrow G$ coincides with $cX \xrightarrow{cf} cNG \cong G$. Now the following diagram

$$\begin{tikzcd}
X \arrow[r, "f"] \arrow[d, "\eta"'] & NG \arrow[d, "\cong"] \\
NcX \arrow[r, "Ncf"'] & NcNG
\end{tikzcd}$$

is commutative, then Ncf is a weak equivalence since η is a weak equivalence from Corollary 7.10. Thus cf is a weak equivalence in $\mathbf{Cat}_1$. Conversely, it is obvious that f is a weak equivalence in $\mathbf{SSets}_1$ if $f^\sharp$ is a weak equivalence in $\mathbf{Cat}_1$. ☐

References

[And78] D. W. Anderson. Fibrations and geometric realizations. Bull. Amer. Math. Soc., 84(5):765-788, 1978.

[CGT04] C.Casacuberta and Marek Golasinski and Andrew Tonks. Homotopy localizations of groupoids. Forum Math., 18, 2006.

[CM06] Claude Cibils and Eduardo N. Marcos. Skew category, Galois covering and smash product of a k-category. Proc. Amer. Math. Soc., 134(1):39-50 (electronic), 2006, arXiv:math/0312214.

[CRS07] Claude Cibils, Maria Julia Redondo, and Andrea Solotar. The Intrinsic Fundamental Group of a Linear Category, arXiv:0706.2491.

[DK80] W. G. Dwyer and D. M. Kan. Calculating simplicial localizations. J. Pure Appl. Algebra, 18(1):17-35, 1980.

[DP95] Carmen Elvira-Donazar and Luis-Javier Hernandez-Paricio. Closed model categories for the n-type of spaces and simplicial sets. Math. Proc. Camb. Phil. Soc. 118:93-103, 1995

[Gab81] Gabriel. P. The universal cover of a representation-finite algebras. volume 903 of Lecture Notes in Math., pages 68-105. Springer, Berlin, 1981

[GZ67] P. Gabriel and M. Zisman. Calculus of fractions and homotopy theory. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 35. Springer-Verlag New York, Inc., New York, 1967.

[Hir03] Philip S. Hirschhorn. Model categories and their localizations, volume 99 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2003.

[Hov99] Mark Hovey. Model categories, volume 63 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 1999.

[Joy08] André Joyal. Notes on Quasi-categories. preprint. 2008.

[JT91] André Joyal and Myles Tierney. Strong stacks and classifying spaces. In Category theory (Como, 1990), volume 1488 of Lecture Notes in Math., pages 213-236. Springer, Berlin, 1991.

[LTW79] D. Latch and R. Thomason and W. Wilson. Simplicial sets from categories. Math. Z. 164:195-214. 1979.

[May92] J. P. May. Simplicial objects in algebraic topology. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1992.

[May99] J. P. May. A concise course in algebraic topology. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1999.

[Min02] Elias Gabriel Minian. Cat as a Λ-cofibration category. J. Pure Appl. Algebra, 167(2-3):301-314, 2002.

[Qui67] Daniel G. Quillen. Homotopical algebra. Lecture Notes in Mathematics, No. 43. Springer-Verlag, Berlin, 1967.

[Qui68] Daniel G. Quillen. The geometric realization of a Kan fibration is a Serre fibration. Proc. Amer. Math. Soc., 19:1499-1500, 1968.

[Rez00] Charles Rezk. A model category for categories. preprint. 2000.

[Str00] N. P. Strickland. K(N)-local duality for finite groups and groupoids. Topology, 39(4):733-772, 2000, arXiv:math.AT/0011109.

[Tho79] R. W. Thomason. Homotopy colimits in the category of small categories. Math. Proc. Comb. Phil. Soc, 85:91-109, 1979.

[Tho80] R. W. Thomason. Cat as a closed model category. Cahiers Topologie Geom. Differentielle, 21(3):305-324, 1980.